

5 TECHNICAL FIELD

10

15

20

Several related U.S. patents address increasing sterol accumulation in higher plants. Those patents include U.S. Patent No. 5,589,619 "Process and Composition for increasing squalene and sterol accumulation in higher plants" (accumulation of squalene in transgenic plants by increasing HMGR activity) and U.S. Patent No. 5,306,862 "Method and composition for increasing sterol accumulation in higher plants" (increasing HMGR activity to increase plant sterol accumulation--including sterol and cycloartenol, which affects insect resistance--in tobacco, tomato, corn, carrot, soybean, cotton, barley, arabidopsis, guayule and petunia; seeds with elevated sterol/cycloartenol, 7S promoter and CaMV promoters), U.S. Patent No. 5,365,017 "Method and composition for increasing sterol accumulation in higher plants" (DNA

[illegible]

WO 9845457 (SMTI, Erg6 from A.t., corn, yeast;
transgenic plants with altered sterol levels_using DNA
encoding an enzyme binding a first sterol and producing

1. *Chlorophyll a* (Chl *a*) is the primary photosynthetic pigment in most algae and higher plants. It is a green pigment that absorbs light energy in the blue and red regions of the visible spectrum. Chl *a* is essential for the light-dependent reactions of photosynthesis, where it converts light energy into chemical energy.

2. *Chlorophyll b* (Chl *b*) is an accessory pigment found in green algae and higher plants. It absorbs light energy in the blue and orange-red regions of the visible spectrum. Chl *b* transfers the absorbed energy to Chl *a* for use in photosynthesis.


3. *Carotenoids* are a group of pigments that include carotenes and xanthophylls. They absorb light energy in the blue and green regions of the visible spectrum. Carotenoids transfer energy to Chl *a* and also play a role in protecting the photosynthetic apparatus from damage by excess light energy.

4. *Phycobilins* are water-soluble pigments found in cyanobacteria and red algae. They absorb light energy in the blue and green regions of the visible spectrum. Phycobilins transfer energy to Chl *a* and are involved in the light-harvesting process.

5. *Phaeophytins* are pigments found in brown algae and some other groups. They are derived from Chl *a* and absorb light energy in the blue and green regions of the visible spectrum. Phaeophytins transfer energy to Chl *a* and are involved in the light-harvesting process.

A key enzyme in sterol biosynthesis is 3-hydroxy-3-methylglutaryl-Coenzyme A reductase (HMG-CoA reductase or HMGR). Schaller et al. (Plant Physiol. 109: 761-770, 1995) found that over-expression of rubber HMGR (*hmg1*) genomic DNA in tobacco leads to the overproduction of sterol end-products (sitosterol, campesterol and stigmasterol) up to 6-fold in leaves. Further, the excess sterol was stored as steryl-esters in lipid bodies. HMGR activity was increased by 4- to 8-fold.

Sterols are derivatives of a fused, reduced ring system, cyclopenta-[a]-phenanthrene, comprising three fused cyclohexane rings (A, B, and C) in a phenanthrene arrangement, and a terminal cyclopentane ring (D) having the formula (I) and carbon atom position numbering shown below:



where R is an 8 to 10 carbon-atom side chain.

5 In plants, squalene is converted to squalene epoxide, which is then cyclized to form cycloartenol (4,4,14 α -trimethyl-9 β ,19-cyclo-5 α -cholest-24-en-3 β -ol). Cycloartenol has two methyl groups at position 4, a methyl group at position 14, a methylene bridge between
10 the carbon atoms at positions 9 and 19 that forms a disubstituted cyclopropyl group at those positions, and includes an 8-carbon sidechain of the formula:
 $\text{CH}_3\text{CH}(\text{CH}_2)_2\text{CH}=\text{C}(\text{CH}_3)_2$. Squalene epoxide can
alternatively be converted into pentacyclic sterols,
15 containing five instead of four rings. Exemplary pentacyclic sterols include the phytoalexins and saponins.

Being one of the first sterols in the higher plant biosynthetic pathway, cycloartenol serves as a
20 precursor for the production of numerous other sterols. In normal plants, cycloartenol is converted to predominantly 24-methylene cycloartenol (4,4,14 α -dimethyl-9 β ,19-cyclo-22,23-dihydro-ergosta-24(28)-en-3 β -ol), cycloeucalenol, (4,14 α -trimethyl-9 β ,19 cyclo-5 α -
25 ergosta-24(28)-en-3 β -ol), isofucosterol (5 α -stigmasta-5-24(28)-dien-3 β -ol), sitosterol (5 α -stigmasta-5-en-3 β -ol), stigmasterol-(stigmasta-5,-22-dien-3 β -ol),

09885729.062004
F00290 E2258860

[illegible]

5

15

25

30

[illegible]

10

15

20

25

30

U.S. Patent No. 5,668,001 teaches a recombinant avian HMG-CoA synthase preparation useful for evaluating drugs that inhibit cholesterol biosynthesis.

JP 09121863 discloses a plant with increased 3-hydroxy-3-methyl glutaryl coenzyme A reductase (HMGR) activity as a result of increasing the expression of a mutant protein kinase gene that regulates expression of the HMGR gene. The increased HMGR activity increases squalene, zymosterol, cholesta-7,24-dienol and cholest-5,7,24-trenol accumulation in yeast with ERG5 and ERG6 mutants.

EP 480730 "Plant-sterol accumulation and pest resistance-by increasing copy number of 3-hydroxy-3-methyl glutaryl coenzyme-A reductase gene in tobacco, tomato and corn

WO 9913086 "Human Delta 7-sterol reductase polypeptide-useful for diagnosis or treatment of genetic defects e.g. hereditary Smith-Lemli-Opitz syndrome" teaches making and using the recombinant polypeptide with humans.

Chappell et al. U.S. Patent No. 5,589,619 teaches that transformation of higher plants with truncated HMG-CoA reductase enhanced the production of squalene, cycloartenol and certain sterols, particularly compounds having unsaturations at the 5-position. Several intermediate sterols as are shown in Fig. 1 were also produced. It would be beneficial if the production of sitosterol and stigmasterol could be enhanced while lessening the accumulation of the intermediate sterols. The present invention provides avenues for enhancing production of sitosterol and stigmasterol and lessening the accumulation of the intermediate sterols.

Gonzalez et al. (Abstract of poster at Third
Terpnet Meeting of the European Network on Plant
Isoprenoids, May 29-30, 1997, Poitiers, France) over-
expressed the *Arabidopsis* HMGR cDNA (*hmg1* and *hmg2*) and
5 found sterol overproduction with *hmg1* only. They used
two forms of the *hmg1* gene, a full-length form and a
truncated form containing only the catalytic domain.
HMGRs have three domains, an N-terminal membrane
spanning domain, a short linker domain, and a C-
10 terminal catalytic domain. In this case the transgenic
plants were also *Arabidopsis*. The difference between
the full-length and truncated forms was a greater
accumulation of pathway intermediates in the case of
the truncated form. More importantly, the intermediates
15 demonstrated as accumulating were cycloartenol, 24-
methylenecycloartanol and obtusifolliol.

Finally, US patents 5,365,017 and 5,306,862, both
assigned to Amoco Corp., disclose a method for
increasing sterol accumulation in plants by increasing
20 the copy number of a gene having HMG-CoA reductase
activity. These inventions disclose a method using
hamster truncated HMGR that consisted of the catalytic
domain and the linker domain. According to the claims
the linker domain was essential for activity. They also
25 demonstrated a greater accumulation of pathway
intermediates such as cycloartenol.

BRIEF SUMMARY

The present invention relates to transgenic plants
30 and their progeny having improved nutritional
characteristics. More particularly, the present
invention relates to transgenic plants and their
progeny, the storage organs (e.g. seed, fruit and
vegetable parts) of which contain modified levels of

steroid compounds, such as (i) elevated levels of beneficial phytosterols (e.g., sitosterol), phytostanols (e.g., sitostanol), and esters thereof, relative to an otherwise identical plant transformed only with a truncated HMG-CoA reductase gene or a wild-type plant, and (ii) reduced levels of steroid pathway intermediate compounds (e.g. one or more of squalene, cycloartenol, 24-methylene cycloartenol, obtusifoliol, stigmasta-7-enol and campesterol) in their storage organs relative to an otherwise identical transgenic plant transformed only with a truncated HMG-CoA reductase gene. Nucleic acid sequences encoding enzymes that affect the biosynthesis and accumulation of steroid compounds in plants (HMG-CoA reductase and a steroid pathway enzyme), and methods for using these sequences to produce such transgenic plants, are also provided. These methods comprise, for example, introducing into cells nucleic acid sequences encoding enzymes that affect the levels of accumulated steroid pathway end products.

The present invention contemplates a recombinant construct or a recombinant vector that contains 2 DNA sequences. The first encodes a polypeptide exhibiting 3-hydroxy-3-methylglutaryl-Coenzyme A (HMG-CoA) reductase activity. The second DNA sequence encodes a polypeptide exhibiting the activity of another steroid pathway enzyme. Each polypeptide-encoding DNA sequence is operably linked in the 5' to 3' direction to a promoter and a transcription termination signal sequence independent of the other sequence. The promoter is located upstream and the termination sequence downstream of each polypeptide-encoding DNA sequence. The second DNA sequence encoding a steroid pathway enzyme can code for a squalene epoxidase

enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifolius C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, or a sterol methyl transferase II enzyme. It is contemplated that
5 HMG-CoA reductase and the steroid pathway enzyme activity comes from a mutant or truncated form of those enzymes, such as a truncated HMG-CoA reductase lacking the transmembrane region while retaining a functional catalytic domain. Examples of such preferred HMG CoA
10 reductases include the truncated rubber and *Arabidopsis* HMG CoA reductases disclosed herein.

Preferably, the regulatory function of a promoter is substantially unaffected by cellular levels of squalene such as the CaMV 35S promoter. In one aspect,
15 a promoter is seed-specific. In another aspect, a promoter is derived from a species in a different order from a host cell. In another aspect, the HMG-CoA reductase or steroid pathway enzymes is from a species in a different order from the order that of the host
20 cell. The invention contemplates a construct or recombinant vector having more than one DNA sequence encoding a steroid pathway enzyme that do not have to be under the control of the same promoter. Preferably, a recombinant vector is a plant expression vector.

25 In another aspect of the invention, a transformed host cell comprises a recombinant construct or vector as described above. Preferably, a host cell is plant cell, preferably that plant cell is from canola, soybean, corn, maize, tobacco, cotton, rape, tomato or
30 alfalfa. The invention contemplates a host cell in a cell culture, plants derived from transformed host cells, and storage organs (seeds, fruits and vegetable parts) from transgenic plants.

In addition to contemplating transgenic plants and seeds, the invention contemplates transgenic plant seeds capable of germinating into a transgenic plant and mutants, recombinants, genetically engineered
5 derivatives thereof and hybrids derived therefrom. The plant over-accumulates steroid pathway products relative to a native, non-transgenic plant of the same strain, wherein said mutants, recombinants, genetically engineered derivatives thereof and hybrids derived
10 therefrom maintain the ability to overaccumulate steroid pathway products.

The invention contemplates a process of increasing the formation of steroid pathway products in a transformed host cell as compared to an otherwise
15 identical non-transformed host cell. Contemplated processes use the described recombinant constructs and vectors to transform host cells, then growing the host cells or regenerating transgenic plants therefrom.

In one aspect of the invention, the genome of a
20 contemplated plant, its progeny, seeds or cell culture, comprises introduced DNA encoding an HMG-CoA reductase activity and introduced DNA encoding a steroid pathway enzyme that is a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase
25 enzyme, a obtusifoliol C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, or a sterol methyl transferase II enzyme. The storage organs of such a plant contain an elevated level of total accumulated sterol, compared to storage organs of an otherwise
30 identical plant, the genome of which does not comprise said introduced DNA. Further, the storage organs of the plant contain a reduced level of squalene, cycloartenol, 24-methylene cycloartenol, obtusifoliol,

09085723.062004
T00290"02258860

stigmasta-7-enol, or campesterol compared to the seeds of an otherwise identical plant or a plant comprising an introduced DNA encoding an HMG-CoA reductase enzyme.

The invention contemplates a method of producing a plant that accumulates an elevated level of sterol pathway products compared to a corresponding plant comprising no introduced DNA encoding a peptide, polypeptide, or protein that affects the biosynthesis and accumulation of a sterol pathway product, comprising sexually crossing plants to arrive at a plant comprising nucleic acid encoding an HMG CoA reductase and a steroid pathway enzyme, including crosses with a nurse cultivar. The plants, including apomictic plants, uniform populations of the plants and their seeds and parts other than seeds are contemplated.

Another aspect of the invention is oils containing at least one sterol pathway product, extracted from the seeds of a contemplated plant. Preferably sitosterol, at least one sitosterol ester, or mixtures thereof, comprise at least about 57% by weight of the total sterol compounds of a contemplated oil. Preferably sitosterol, that at least one sitosterol ester, or mixtures thereof, comprise at least about 0.08% of the dry weight of a contemplated seed. Preferably, the oil has a reduced amount of squalene, cycloartenol, 24-methylene cycloartenol, obtusifoliol, stigmasta-7-enol, campesterol, or combinations thereof, compared to oil from a corresponding transgenic plant that does not contain introduced DNA encoding a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifoliol C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, a sterol methyl

transferase II enzyme, or mixture thereof; wherein the reduction is in the range of from about 10% to about 100%.

Sitosterol ester compositions derived from
5 transgenic plants of the present invention, their progeny or their seeds are also contemplated, preferably wherein an esterifying fatty acid has 2 to 22 carbon atoms in the main chain.

A further aspect of the invention is cholesterol-
10 lowering compositions comprising contemplated oils and sitosterol ester compositions. Another further aspect of the invention is foods, food ingredients, or food compositions comprising contemplated oils.

Still further, the invention contemplates
15 pharmaceutical compositions comprising a cholesterol-lowering effective amount of a contemplated oil, and a pharmaceutically acceptable carrier, excipient, or diluent.

A method of lowering the plasma concentration of
20 low density lipoprotein cholesterol is contemplated, comprising orally administering to a human or animal subject an effective amount of an above composition. Also contemplated is a method of treating or preventing an elevated plasma concentration of low-density
25 lipoprotein cholesterol, comprising orally administering to a human or animal subject an effective amount of a contemplated composition.

A related aspect of the invention is a method of making a food additive composition, comprising
30 obtaining oil containing a sterol pathway product compound from seed of a contemplated transgenic plant and mixing the oil with an edible solubilizing agent, an effective amount of a dispersant, and optionally, an effective amount of an antioxidant.

09885723.062001
T00250.22258850

Novel forms of two sterol pathway enzymes and the nucleic acids that encode them are disclosed: an *Arabidopsis* enzyme having nucleic acid similarity to a squalene epoxidase, and an *Arabidopsis* enzyme having

5 nucleic acid similarity to an obtusifoliol C14 α -demethylase enzyme. Thus, the invention contemplates an isolated DNA molecule having a nucleotide sequence of disclosure SEQ ID NO: 4, 6, 8, 10, 14, 15, 17 or the complements thereof. Also contemplated is a nucleotide

10 sequence that hybridizes to the nucleotide sequence of SEQ ID NO:4, 6, 8, 10, 14, 15, 17 or their complements under a wash stringency equivalent to 0.5X SSC to 2X SSC, 0.1% SDS, at 55-65°C, and that encode a polypeptide having squalene epoxidase or obtusifoliol

15 C14 α -demethylase enzymatic activity. Preferably, that enzymatic activity is substantially similar to that of a disclosed squalene epoxidase or obtusifoliol C14 α -demethylase, respectively. By substantially similar is meant having enzymatic activity differing from that

20 of the disclosed enzymes by about 30% or less, preferably by about 20% or less, and more preferably by about 10% or less when assayed by standard enzymatic assays. Also contemplated is a nucleotide sequence encoding the same genetic information as said

25 nucleotide sequence of SEQ ID NO: 4, 6, 8, 10, 14, 15, 17 or their complements or that hybridize as described above, but which is degenerate in accordance with the degeneracy of the genetic code. Recombinant constructs, vectors and transformed host cells

30 comprising the novel isolated and purified nucleic acid sequences are also contemplated. In one embodiment, the vector is a plant vector and the host cell is a plant cell. Methods of producing the disclosed

squalene epoxidase or obtusifoliol C14 α -demethylase enzymes are also contemplated comprising culturing a transformed host cell for a time and under conditions conducive to the production of the squalene epoxidase or obtusifoliol C14 α -demethylase enzyme, and recovering the produced squalene epoxidase or obtusifoliol C14 α -demethylase enzyme.

Yet another aspect provides any of the above described transformed host cells, further comprising a recombinant construct or expression vector encoding a tocopheral synthesis pathway enzyme, and in particular, S-adenosylmethionine-dependent α -tocopherol methyltransferase. Also included are plants, seeds and storage organs comprising the transformed host cells.

Another aspect provides, a process of increasing the formation of steroid pathway products and tocopherols in a transformed host cell as compared to an otherwise identical non-transformed host cell comprising (1) transforming a host cell with a recombinant vector comprising (a) as operably linked components in the 5' to 3' direction, a promoter, a DNA sequence encoding a first polypeptide having 3-hydroxy-3-methylglutaryl-Coenzyme A reductase enzyme activity, and a transcription termination signal sequence; and (b) as operably linked components in the 5' to 3' direction, a promoter, a DNA sequence encoding at least one polypeptide having steroid pathway enzyme activity selected from the group consisting of squalene epoxidase enzyme activity, sterol methyl transferase I enzyme activity, sterol C4-demethylase enzyme activity, obtusifoliol C14 α -demethylase enzyme activity, sterol C5-desaturase enzyme activity, and sterol methyl transferase II enzyme activity, and a transcription

termination signal sequence; (2) transforming the host cell of (1) with a recombinant vector comprising as operably linked components, a promoter, a DNA sequence encoding a tocopherol synthesis pathway enzyme, and a transcription termination sequence; and (3) regenerating said transformed plant cell into said transgenic plant.

10 BRIEF DESCRIPTION OF THE DRAWINGS

These and other features, aspects, and advantages of the present invention will become better understood with regard to the following description, appended claims and accompanying figures where:

15 Fig. 1 is an abbreviated version of a plant steroid compound biosynthetic pathway that shows the enzymes affecting steroid compound biosynthesis and accumulation. These include: HMG-CoA reductase, squalene epoxidase, sterol methyl transferase I, sterol C4-demethylase, obtusifoliol C14 α -demethylase, sterol C5 desaturase and sterol methyl transferase II.

Fig. 2 depicts the forms of *Arabidopsis* and rubber HMGR1 tested in *Arabidopsis* and yeast to compare expression, activity and sterol production.

25 Fig. 3 is a map showing the structure of construct pMON29920. pMON29920 is a binary transformation vector with P-7S/E9 3' cassette and the KAN gene flanked by the two borders where P-7S is the promoter of alpha' beta conglycinin protein from soybean, E9 3' is the 3' end of pea rbc E9 gene and KAN is the coding sequence for NPTII that confers resistance to kanamycin. The NPTII gene is driven by the 35S promoter from cauliflower mosaic virus. Spc.Str is the coding region

30

Table 1. *Continued*

Fig. 5 is a map showing the structure of construct pMON23616. pMON23616 is a plant expression vector containing P-NOS/ORF-7/KAN/NOS-3' cassette. P-NOS: NOS promoter from *Agrobacterium tumefaciens* pTiT37; ORF-7: a short open reading frame that attenuates expression of KAN in plants; KAN: coding sequence of NPTII gene that confers resistance to kanamycin and neomycin; ble: confers resistance to bleomycin; NOS 3': 3' termination end of nopaline synthase coding region; Left Border: Octapine left border from Octapine Ti plasmid pTiA6; ori-V: the vegetative origin of replication; rop:

coding region for repressor of primer; Spc/Str: coding region for Tn7 adenylyltransferase conferring resistance to spectinomycin and streptomycin.

Fig. 6 is a map showing the structure of construct pMON43818. pMON43818 is a recombinant binary vector carrying the gene encoding rubber hydroxymethyl glutaryl CoA reductase1 (HMGR1) in sense orientation driven by the soybean alpha' beta conglycinin promoter. P-NOS: nopaline synthase gene promoter; kan: coding region for neomycin phospho transferase protein to confer resistance to kanamycin; NOS 3': 3' termination end of nopaline synthase coding region; Soy Alpha' Beta Conglycinin: 7S alpha' beta conglycinin gene promoter from soybean; Rubber HMGR1 gene: coding sequence for HMGR1 gene from *Hevea brasiliensis*; E9 3': 3' end of pea rbcS E9 gene; Left border: octopine left border, sequence essential for transfer of T-DNA into *Agrobacterium*; ori-V: plasmid origin of replication in *Agrobacterium*; rop: coding sequence for repressor of primer; Ori-322: origin of replication in *E.coli*; Spc/Str: coding region for Tn7 adenyllyltransferase (AAD(3")) conferring resistance to spectinomycin and streptomycin; Right Border: right border sequence of T-DNA essential for integration into *Agrobacterium*.

Figure 7 is a map showing the structure of construct pMON43052. pMON43052 is a recombinant shuttle vector, carrying the cDNA fragment encoding the catalytic domain of *Arabidopsis* HMGR1 in sense orientation driven by the soybean alpha' beta conglycinin promoter. P-NOS: nopaline synthase gene promoter; kan: coding region for neomycin phosphotransferase protein to confer resistance to kanamycin; NOS 3': 3' termination end of nopaline synthase coding region; Soy Alpha' Beta Conglycinin: 7S

10 adenylyltransferase (AAD(3")) conferring resistance to
spectinomycin and streptomycin; Right Border: right
border sequence of T-DNA essential for integration into
Agrobacterium.

20 end of nopaline synthase coding region; Left border:
octopine left border sequence essential for transfer of
T-DNA into *Agrobacterium*; ori-V: plasmid origin of
replication in *Agrobacterium*; rop: coding sequence for
repressor of primer; ori-322: origin of replication in
25 *E.coli*; Spc/Str: coding region for Tn7
adenylyltransferase (AAD(3")) conferring resistance to
spectinomycin and streptomycin; Right Border: right
border sequence of T-DNA essential for integration into
Agrobacterium.

Figure 9 is a map showing the structure of construct pMON43057. pMON43057 is a recombinant binary vector for *Agrobacterium* mediated transformation of soybean, carrying the gene cassette for expressing catalytic domain of HMGR1 from *Arabidopsis thaliana*.

The catalytic domain of the HMGR1 cDNA is driven by soybean 7S alpha' beta conglycinin promoter. P-NOS: nopaline synthase gene promoter; kan: coding region for neomycin phosphotransferase protein to confer

5 resistance to kanamycin; NOS 3': 3' termination end of nopaline synthase coding region; Left border: octopine left border sequence essential for transfer of T-DNA into *Agrobacterium*; ori-V: plasmid origin of replication in *Agrobacterium*; rop: coding sequence for repressor of primer; ori-322: origin of replication in *E.coli*; Spc/Str: coding region for Tn7

10 adenylyltransferase (AAD(3")) conferring resistance to spectinomycin and streptomycin; Right Border: right border sequence essential for transfer of T-DNA into *Agrobacterium*; Soy Alpha' Beta Conglycinin: soybean 7S

15 alpha' beta conglycinin gene promoter; *Arabidopsis* HMGR catalytic domain: coding sequence for *Arabidopsis* HMGR1 catalytic domain; E9 3': 3' end of pea rbcS E9 gene.

20 Figure 10 is a map showing the structure of construct pMON43058. pMON43058 is a recombinant binary vector for *Agrobacterium*-mediated soybean transformation, carrying gene expression cassettes for catalytic domain of HMGR1 from *Arabidopsis thaliana* and

25 SMTII from *Arabidopsis thaliana*. P-NOS: nopaline synthase gene promoter; kan: coding region for neomycin phosphotransferase protein to confer resistance to kanamycin; NOS 3': 3' termination end of nopaline synthase coding region; Left border: octopine left

30 border sequence essential for transfer of T-DNA into *Agrobacterium*; ori-V: plasmid origin of replication in *Agrobacterium*; rop: coding sequence for repressor of primer; ori-322: origin of replication in *E.coli*; Spc/Str: coding region for Tn7 adenylyltransferase

09885723.062001
"000290" 02/08/95

(AAD(3")) conferring resistance to spectinomycin and streptomycin; Right Border: right border sequence essential for transfer of T-DNA into *Agrobacterium*; Soy Alpha' Beta Conglycinin: 7S alpha' beta conglycinin gene promoter from soybean; *Arabidopsis* HMGR catalytic domain: sequence encoding the catalytic domain of *Arabidopsis* HMGR1; E9 3': 3' end of pea rbcS E9 gene; Soy Alpha' Beta Conglycinin: soybean 7S alpha'beta conglycinin gene promoter; *Arabidopsis* SMT2: cDNA encoding sterol methyl transferase II enzyme from *Arabidopsis thaliana* (accession no: X89867); NOS 3': 3' termination end of nopaline synthase coding region.

Fig. 11 is profile (histogram) of the sterol composition of R1 transgenic soybean seeds when *Arabidopsis* truncated HMGR (catalytic domain without linker) was overexpressed using seed-specific 7S promoter (data from pMON43057:p7S::At HMGR truncated).

Fig. 12 is a profile (histogram) of the sterol composition of R1 transgenic soybean seeds when *Arabidopsis* truncated HMGR (catalytic domain without linker) and *Arabidopsis* SMTII were overexpressed (data from pMON43058:p7S::At HMGR truncated and p7S::At SMTII). The expression of the genes is controlled by the seed-specific 7S promoter.

Figure 13 is a map showing the structure of construct pMON53733. pMON53733 is a recombinant binary vector carrying the cDNA encoding full-length form of *Arabidopsis* hydroxymethyl glutaryl CoA reductase1 (HMGR1) in sense orientation driven by the enhanced cauliflower mosaic virus 35S promoter. P-35S: 35S promoter from cauliflower mosaic virus; kan: confers resistance to neomycin and kanamycin; NOS 3': 3' termination end of nopaline synthase coding region; Left border: octopine left border, sequence essential

09085723.062004
T00290" 02450000

for transfer of T-DNA into *Agrobacterium*; ori-V:
 plasmid origin of replication in *Agrobacterium*; rop:
 coding sequence for repressor of primer; ori-322:
 origin of replication in *E.coli*; Spc/Str: coding region
 5 for Tn7 adenyltransferase (AAD(3")) conferring
 resistance to spectinomycin and streptomycin; Right
 Border: right border sequence of T-DNA essential for
 integration into *Agrobacterium*; P-e35S: enhanced
 cauliflower mosaic virus promoter; *Arabidopsis* HMGR1:
 10 cDNA sequence encoding full-length form of *Arabidopsis*
 HMGR1; E9 3': 3' end of pea rbcS E9 gene.

Figure 14 is a map showing the structure of
 construct pMON53734. pMON53734 is a recombinant binary
 vector carrying the cDNA encoding catalytic domain with
 15 linker region of *Arabidopsis* hydroxymethyl glutaryl CoA
 reductase1 (HMGR1) in sense orientation driven by the
 enhanced cauliflower mosaic virus 35S promoter. P-35S:
 35S promoter from cauliflower mosaic virus; kan:
 confers resistance to neomycin and kanamycin; NOS 3':
 20 3' termination end of nopaline synthase coding region;
 Left border: octopine left border, sequence essential
 for transfer of T-DNA into *Agrobacterium*; ori-V:
 plasmid origin of replication in *Agrobacterium*; rop:
 coding sequence for repressor of primer; ori-322:
 25 origin of replication in *E.coli*; Spc/Str: coding region
 for Tn7 adenyltransferase (AAD(3")) conferring
 resistance to spectinomycin and streptomycin; Right
 Border: right border sequence of T-DNA essential for
 integration into *Agrobacterium*; P-e35S: enhanced
 30 cauliflower mosaic virus promoter; *Arabidopsis* tHMGR1:
 cDNA sequence encoding catalytic domain with linker
 region of *Arabidopsis* HMGR1; E9 3': 3' end of pea rbcS
 E9 gene.

0985723.062004
T00290"0258860

5

25

30

coding sequence for repressor of primer; ori-322: origin of replication in *E.coli*; Spc/Str: coding region for Tn7 adenylyltransferase (AAD(3")) conferring resistance to spectinomycin and streptomycin; Right Border: right border sequence of T-DNA essential for integration into *Agrobacterium*; P-e35S: enhanced cauliflower mosaic virus promoter; Hevea HMGR1: cDNA sequence encoding full-length form of rubber HMGR1; E9 3': 3' end of pea rbcS E9 gene.

Figure 17 is a map showing the structures of construct pMON53737. pMON53737 is a recombinant binary vector carrying the cDNA encoding catalytic domain with linker region of rubber (*Hevea brasiliensis*) hydroxymethyl glutaryl CoA reductase1 (HMGR1) in sense orientation_driven by the enhanced cauliflower mosaic virus 35S promoter. P-35S: 35S promoter from cauliflower mosaic virus; kan: confers resistance to neomycin and kanamycin; NOS 3': 3' termination end of nopaline synthase coding region; Left border: octopine left border, sequence essential for transfer of T-DNA into *Agrobacterium*; ori-V: plasmid origin of replication in *Agrobacterium*; rop: coding sequence for repressor of primer; ori-322: origin of replication in *E.coli*; Spc/Str: coding region for Tn7 adenyllyltransferase (AAD(3")) conferring resistance to spectinomycin and streptomycin; Right Border: right border sequence of T-DNA essential for integration into *Agrobacterium*; P-e35S: enhanced cauliflower mosaic virus promoter; rubber tHMGR1: cDNA sequence encoding catalytic domain with linker region of rubber HMGR1; E9 3': 3' end of pea rbcS E9 gene.

Figure 18 is a map showing the structure of construct pMON53738. pMON53738 is a recombinant binary vector carrying the cDNA encoding mutant form of rubber

(*Hevea brasiliensis*) hydroxymethyl glutaryl CoA reductase1 (HMGR1) in sense orientation driven by the enhanced cauliflower mosaic virus 35S promoter. In the mutant rubber HMGR1 the putative phosphorylation site, the serine amino acid residue at position 566 is changed to alanine amino acid residue (SEQ ID 23). P-35S: 35S promoter from cauliflower mosaic virus; kan: confers resistance to neomycin and kanamycin; NOS 3': 3' termination end of nopaline synthase coding region; Left border: octopine left border, sequence essential for transfer of T-DNA into *Agrobacterium*; ori-V: plasmid origin of replication in *Agrobacterium*; rop: coding sequence for repressor of primer; ori-322: origin of replication in *E.coli*; Spc/Str: coding region for Tn7 adenyltransferase (AAD(3")) conferring resistance to spectinomycin and streptomycin; Right Border: right border sequence of T-DNA essential for integration into *Agrobacterium*; P-e35S: enhanced cauliflower mosaic virus promoter; rubber tHMGR1 Ala 566: cDNA sequence encoding catalytic domain with linker region of rubber HMGR1 in which serine amino acid residue at position 566 is changed to alanine amino acid residue using site directed mutagenesis; E9 3': 3' end of pea rbcS E9 gene.

Figure 19 is a map showing the structure of construct pMON53739. pMON53739 is a recombinant binary vector carrying the cDNA encoding mutant form of rubber (*Hevea brasiliensis*) hydroxymethyl glutaryl CoA reductase1 (HMGR1) in sense orientation driven by the enhanced cauliflower mosaic virus 35S promoter. In the mutant rubber HMGR1 the putative phosphorylation site, the serine amino acid residue at position 567 is changed to alanine amino acid residue (SEQ ID 24). P-35S: 35S promoter from cauliflower mosaic virus; kan:

confers resistance to neomycin and kanamycin; NOS 3':
 3' termination end of nopaline synthase coding region;
 Left border: octopine left border, sequence essential
 for transfer of T-DNA into *Agrobacterium*; ori-V:
 5 plasmid origin of replication in *Agrobacterium*; rop:
 coding sequence for repressor of primer; ori-322:
 origin of replication in *E.coli*; Spc/Str: coding region
 for Tn7 adenyltransferase (AAD(3")) conferring
 resistance to spectinomycin and streptomycin; Right
 10 Border: right border sequence of T-DNA essential for
 integration into *Agrobacterium*; P-e35S: enhanced
 cauliflower mosaic virus promoter; rubber tHMGR1 Ala
 567: cDNA sequence encoding catalytic domain with
 linker region of rubber HMGR1 in which serine amino
 15 acid residue at position 567 is changed to alanine
 amino acid residue using site directed mutagenesis; E9
 3': 3' end of pea rbcS E9 gene.

Figure 20 is a map showing the structure of
 construct pMON53740. pMON53740 is a recombinant binary
 20 vector carrying the cDNA encoding catalytic domain
 without linker region of rubber (*Hevea brasiliensis*)
 hydroxymethyl glutaryl CoA reductase1 (HMGR1) in sense
 orientation driven by the enhanced cauliflower mosaic
 virus 35S promoter. P-35S: 35S promoter from
 25 cauliflower mosaic virus; kan: confers resistance to
 neomycin and kanamycin; NOS 3': 3' termination end of
 nopaline synthase coding region; Left border: octopine
 left border, sequence essential for transfer of T-DNA
 into *Agrobacterium*; ori-V: plasmid origin of
 30 replication in *Agrobacterium*; rop: coding sequence for
 repressor of primer; ori-322: origin of replication in
E.coli; Spc/Str: coding region for Tn7
 adenyltransferase (AAD(3")) conferring resistance to
 spectinomycin and streptomycin; Right Border: right

098572.0001
T00290" 6253860

5 E9 3': 3' end of pea rbcS E9 gene.

10 to control plants.

15 compared to control plants.

20 to control plants.

25 to control plants.

30 to control plants.

Fig. 26 is a graph comparing the sitostanol content in micrograms of steroid compound per gram of seeds analyzed in transgenic *Arabidopsis* plants

transformed with pMON53733 through pMON53740 compared to control plants.

Fig. 27 is a sterol profile (histogram) of transgenic *Arabidopsis* harboring different forms of rubber HMGR.

Fig. 28 is a graph of the squalene, zymosterol and erogosterol content in micrograms of sterol per milligram of cell dry weight from HMGR constructs in yeast HMGR1 knockout mutants for constructs having full length and truncated HMG CoA reductase (HMGR) sequences. The truncated sequences contain substantial portions of the catalytic region but lack the linker region and the transmembrane region of HMGR. These sequences are derived from *Arabidopsis* and rubber plants.

Fig. 29 is a map showing the structure of construct pMON43842. pMON43842 is a yeast expression vector carrying cDNA encoding *Arabidopsis* putative obtusifoliol C14 α -demethylase (AC002329) in sense orientation driven by the p423Gall1 promoter. Sc.His3: HIS3 region from *Saccharomyces cerevisiae* encoding imidazoleglycerol-phosphate dehydratase for histidine synthesis; Ori-f1: bacteriophage f1 origin of replication; LAC: contains partial lacI coding sequence, promoter Plac, promoter Pt7, promoter Pt3, KS polylinker, and partial lacZ coding sequence; lacZ: partial coding sequence for beta-d-galactosidase or lacZ protein; T-Sc.Cycl: a terminator from Cycl- iso-1-cytochrome c from *Saccharomyces cerevisiae* to terminate transcription; obtus. C14 α .demethylase (AC002329): cDNA encoding *Arabidopsis* putative obtusifoliol C14 α -demethylase; P-Sc.Gall1: a promoter from Gall1- galactokinase of *Saccharomyces cerevisiae* to

098572.062001
T00290" E228860

direct expression with galactose induction; LacZ-
alpha: partial coding sequence for beta-d-galactosidase
or lacZ protein; Ori-pUC: minimum sequence required
for a functional origin of replication, sequence
5 downstream of this region is known to affect copy
number when expressed in bacteria; AMP: contains the
P3 promoter and the beta-lactamase coding sequence,
conferring resistance to ampicillin, penicillin, and
carbenicillin; Sc.2micron: 2 micron origin of
10 replication.

Fig. 30 is a map showing the structure of
construct pMON43843. pMON43843 is a yeast expression
vector carrying cDNA encoding *Arabidopsis* putative
squalene epoxidase 1 (ATA506263) in sense orientation
15 driven by the p423Gal1 promoter. Sc.His3: HIS3 region
from *Saccharomyces cerevisiae* encoding
imidazoleglycerol-phosphate dehydratase for histidine
synthesis; Ori-f1: bacteriophage f1 origin of
replication; LAC: contains partial lacI coding
20 sequence, promoter Plac, promoter Pt7, promoter Pt3, KS
polylinker, and partial lacZ coding sequence; lacZ:
partial coding sequence for beta-d-galactosidase or
lacZ protein; T-Sc.Cyc1: a terminator from Cyc1- iso-
1-cytochrome c from *Saccharomyces cerevisiae* to
25 terminates transcription; Squalene epoxidase 1
(ATA506263): cDNA encoding *Arabidopsis* putative
squalene epoxidase 1 (ATA506263); P-Sc.Gal1: a
promoter from Gal1- galactokinase of *Saccharomyces*
cerevisiae to direct expression with galactose
30 induction; LacZ-alpha: partial coding sequence for
beta-d-galactosidase or lacZ protein; Ori-pUC: minimum
sequence required for a functional origin of
replication, sequence downstream of this region is
known to affect copy number when expressed in bacteria;

D00290" 62458860

AMP: contains the P3 promoter and the beta-lactamase coding sequence, conferring resistance to ampicillin, penicillin, and carbenicillin; Sc.2micron: 2 micron origin of replication.

5 Fig. 31 is a map showing the structure of construct pMON43844. pMON43844 is a yeast expression vector carrying cDNA encoding *Arabidopsis* putative squalene epoxidase 1 (ATA304243) in sense orientation driven by the p423Gal1 promoter. Sc.His3: HIS3 region
10 from *Saccharomyces cerevisiae* encoding imidazoleglycerol-phosphate dehydratase for histidine synthesis; Ori-fl: bacteriophage fl origin of replication; LAC: contains partial lacI coding sequence, promoter Plac, promoter Pt7, promoter Pt3, KS
15 polylinker, and partial lacZ coding sequence; lacZ: partial coding sequence for beta-d-galactosidase or lacZ protein; T-Sc.Cycl: a terminator from Cycl- iso-1-cytochrome c from *Saccharomyces cerevisiae* to terminate transcription; Arab. squalene epoxidase 1
20 (ATA304243): cDNA encoding *Arabidopsis* putative squalene epoxidase 1 (ATA304243); P-Sc.Gal1: a promoter from Gal1- galactokinase of *Saccharomyces cerevisiae* to direct expression with galactose induction; LacZ-alpha: partial coding sequence for
25 beta-d-galactosidase or lacZ protein; Ori-pUC: minimum sequence required for a functional origin of replication, sequence downstream of this region is known to affect copy number when expressed in bacteria; AMP: contains the P3 promoter and the beta-lactamase
30 coding sequence, conferring resistance to ampicillin, penicillin, and carbenicillin; Sc.2micron: 2 micron origin of replication.

Fig. 32 is a comparison of known HMG CoA reductase amino acid sequences. ClustalW alignment of

09885729.0001
"00290"02258860

forty-three non-redundant HMG-CoA reductase sequences to represent archaebacterial, eubacterial, fungal, plant and animal groups. The putative functional domains in the alignment marked as described below are based on the three dimensional structure of *Pseudomonas mevalonii* HMGR (Lawrence et al., 1995, Science 268:1758): boxed-HMGCoA binding domain, light shade-NAD(H) binding domain, underlined consensus-domains involved in catalysis, * underneath consensus and boldface-key histidine residue involved in catalysis. The putative phosphorylation site residues are marked with ‡ and boldface, and are located at the C-terminal region of the protein, adjacent to a highly conserved arginine, marked with † and boldface. Also indicated are the conserved Glu (E), Lys (K), and Asp (D) residues, marked by E, K, and D, respectively. These residues are thought to be critical in catalysis, based on the crystal structure (Tabernero et al., 1999; PNAS 96(13):7167-71).

Appendices A through C show SEQ ID Nos: 1 through 3, respectively. Appendices D through G show SEQ ID Nos 20 thorough 23, respectively.

DETAILED DESCRIPTION

The following detailed description is provided to aid those skilled in the art in practicing the present invention. Even so, this detailed description should not be construed to unduly limit the present invention as modifications and variations in the embodiments discussed herein can be made by those of ordinary skill in the art without departing from the spirit or scope of the present inventive discovery.

All publications, patents, patent applications, databases and other references cited in this

5 incorporated by reference.

10 higher in seeds from these transgenic plants. We also demonstrated a higher accumulation of pathway intermediates in soybean than canola. These results were disclosed in PCT publication WO 00/61771.

15 *Arabidopsis hmg1* without the linker and membrane
spanning domains in *Arabidopsis* and soybean. The
results in *Arabidopsis* were similar to that
demonstrated by Gonzalez et al. (1997) and we compared
the sterol profiles of our transgenic plants with those
20 produced by Gonzalez et al., using our methods to show
they are comparable. We found the same types of pathway
intermediates accumulating. However, in soybean seeds
we have demonstrated the accumulation of squalene to a
very high level (~3mg/g seed which is around 100-fold
25 higher than in nontransgenic controls). This is an
unexpected result not disclosed or suggested in the
prior art. Squalene is a precursor for sterols and in
soybean it appears that there is a "bottleneck" in the
further conversion of this precursor to sterols. Thus,
30 it appears that there could be additional ways of over-
producing sterols in soybean to levels greater than 10-
fold which would include combining a truncated form of
HMGR with other genes coding for enzymes down-stream of
squalene.

This opens the potential to combine other genes such as squalene epoxidase for further enhancing the levels of desirable sterols. Such a combination has not been disclosed or suggested in the prior art.

- 5 Squalene epoxidase catalyzes the addition of oxygen to squalene which is a 30-carbon linear isoprenoid chain thus allowing for cyclization to form cycloartenol. Additional enzymes downstream that can be also be manipulated are sterol methyltransferase 1, C-4
- 10 demethylase, C-14 demethylase, sterol methyltransferase 2, and C-5 desaturase that would all deplete other pathway intermediates shown to accumulate in soybeans. By using such strategies it is possible to convert all of the squalene and other intermediates to end sterols
- 15 such as sitosterol, stigmasterol and campesterol. Thus, sterol level in soybean oils can be elevated from 0.3% up to 3.5%. Expression of the full-length rubber HMGR in soybeans results in a sterol level increase up to 2.7%.
- 20 Enhancement of 3-hydroxy-3-methylglutaryl-CoA reductase (HMG Co-A reductase) activity in certain cells results in increased sterol biosynthesis. See, e.g. Chappell, U.S. Patent No. 5,589,619. The present discovery further contemplates an increase of steroid
- 25 pathway end products such as Δ^5 sterols and their stanol counterparts with a decreased accumulation of certain steroid pathway intermediates by also enhancing various specific steroid pathway enzyme activities, such that more of the steroid pathway intermediate
- 30 compounds are converted to steroid pathway end products.

DNA sequences encoding squalene epoxidases are useful for removal of squalene accumulation, genes

encoding sterol methyl transferase I enzymes are useful for removal of cylcoartenol accumulation, genes encoding sterol C4-demethylase are useful for removal of 24-methylene cycloartenol accumulation, genes
5 encoding obtusifoliol C14 α -demethylases are useful for removal of accumulation of obtusifoliol, genes encoding sterol C5-desaturases are useful for removal of stigmasta-7-enol accumulation, and genes encoding sterol methyl transferase II enzymes are useful for the
10 reduction of accumulated campesterol and concomitant increase of sitosterol.


Levels of sitostanol and sitostanol esters can be elevated further by approximately 2- to 40-fold over the transgenic plants of the art having only added
15 genes for HMG CoA reductase by introducing additional genes encoding one or more of the following sterol pathway enzymes: a squalene epoxidase, a sterol methyl transferase I, a sterol C4-demethylase, an obtusifoliol C14 α -demethylase, a sterol C5-desaturase, a sterol
20 methyl transferase II.

As used herein, the term "structural coding sequence" means a DNA sequence which encodes for a peptide, polypeptide, or protein which may be made by a cell following transcription of the DNA to mRNA,
25 followed by translation to the desired peptide, polypeptide, or protein.

The term "sterol" as applied to plants refers to any chiral tetracyclic isopentenoid which may be formed by cyclization of squalene oxide through the transition
30 state possessing stereochemistry similar to the *trans-syn-trans-anti-trans-anti* configuration, i.e., protosteroid cation, and which retains a polar group at C-3 (hydroxyl or keto), an *all-trans-anti*

stereochemistry in the ring system, and a side-chain
20R-configuration (Parker et al. (1992) *In Nes et al.*,
Eds., *Regulation of Isopentenoid Metabolism*, ACS
Symposium Series No. 497, p. 110; American Chemical
5 Society, Washington, D.C.). The numbering of the
carbon atoms of a representative sterol (cholesterol)
is shown in the following structure (FORMULA II):

As used herein, the term "sterol" refers to
unsaturated hydroxyl group-containing derivatives of a
10 fused, reduced ring system, cyclopenta[α]-phenanthrene,
comprising three fused cyclohexane rings (A, B and C)
in a phenanthrene arrangement, and a terminal
cyclopentane ring (D). The exemplary steroid below
(FORMULA II) illustrates the numbering system employed
15 herein in describing the location of groups and
substituents.



Sterols may or may not contain a C-5 to C-6 double
20 bond, as this is a feature introduced late in the
biosynthetic pathway (note Scheme 1, below). Sterols
contain a C₈-C₁₀ side chain at the C-17 position, as
shown above.

The term "phytosterol," which applies to sterols
25 found uniquely in plants, refers to a sterol containing
a C-5, and in some cases a C-22, double bond.
Phytosterols are further characterized by alkylation of

the C-17 side-chain with a methyl or ethyl substituent at the C-24 position. Major phytosterols include, but are not limited to, sitosterol, stigmasterol, campesterol, brassicasterol, etc. Cholesterol, which
5 lacks a C-24 methyl or ethyl side chain, is found in plants but is not unique thereto, and is not a "phytosterol"

"Phytostanols" are saturated forms of phytosterols wherein the C-5 and, when present, C-22 double bond(s)
10 is(are) reduced, and include, but are not limited to, sitostanol, campestanol, and 22-dihydrobrassicastanol.

"Phytosterol esters" and "phytostanol esters" are further characterized by the presence of a fatty acid or phenolic acid moiety rather than a hydroxyl group at
15 the C-3 position.

The term "steroid compounds" includes sterols, phytosterols, phytosterol esters, phytostanols, and phytostanol esters.

The term "phytosterol compound" refers to at least
20 one phytosterol, at least one phytosterol ester, or a mixture thereof.

The term "phytostanol compound" refers to at least one phytostanol, at least one phytostanol ester, or a mixture thereof.

25 The term "constitutive promoter" refers to a promoter that operates continuously in a cell, and which is not subject to quantitative regulation. The gene with which such a promoter is associated is always "turned on."

30 The terms "seed-specific," "fruit-specific," "plastid-specific," etc., as they apply to promoters refer to preferential or exclusive activity of these promoters in these organs or organelles, respectively. "Preferential expression" refers to promoter activity

09885723.062004
T00290.2228860

[illegible]

As used herein "isolated polynucleotide" means a polynucleotide that is free of one or both of the nucleotide sequences which flank the polynucleotide in the naturally-occurring genome of the organism from which the polynucleotide is derived. The term includes, for example, a polynucleotide or fragment thereof that is incorporated into a vector or expression cassette; into an autonomously replicating plasmid or virus; into the genomic DNA of a prokaryote or eukaryote; or that exists as a separate molecule independent of other polynucleotides. It also includes a recombinant polynucleotide that is part of a hybrid polynucleotide, for example, one encoding a polypeptide sequence.

As used herein "polynucleotide" and "oligonucleotide" are used interchangeably and refer to a polymeric (2 or more monomers) form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. Although nucleotides are usually joined by phosphodiester linkages, the term also includes polymeric nucleotides containing neutral amide backbone linkages composed of aminoethyl glycine units. This term refers only to the primary structure of the molecule. Thus, this term includes double- and single-stranded DNA and RNA. It also includes known types of modifications, for example, labels, methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates,

phosphotriesters, phosphoamidates, carbamates, etc.), those containing pendant moieties, such as, for example, proteins (including for e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide. Polynucleotides include both sense and antisense strands.

The alternative nucleotide sequences described above are considered to possess substantially similar enzymatic activity to that of the polypeptide-encoding polynucleotide sequences of the present invention if they encode polypeptides having enzymatic activity differing from that of any of the polypeptides encoded by the polynucleotide sequences of the present invention by about 30% or less, preferably by about 20% or less, and more preferably by about 10% or less when assayed by standard enzymatic assays.

As used herein "effective amount" is intended to qualify the amount of an agent which will achieve the goal of a lessening in the severity and/or the frequency of incidence of a disease condition or disorder, over no treatment.

The phrase "steroid pathway products" refers to the products of steroid biosynthesis produced by the action of one or more of squalene epoxidase enzyme, sterol methyl transferase I enzyme, sterol C4-demethylase enzyme, obtusifoliol C14 α -demethylase enzyme, sterol C5-desaturase enzyme, and sterol methyl

09885723.062004
T00290" E2458860

transferase II enzyme. Specific examples of steroid pathway products include, but are not limited to, sitosterol, sitostanol, stigmasterol and stigmastanol.

In the context of the present disclosure, a "non-transformed" plant or cell refers to a plant or cells which does not comprise introduced polynucleotides encoding a polypeptide having 3-hydroxy-3-methylglutaryl-Coenzyme A reductase enzyme activity and at least one polypeptide having squalene epoxidase enzyme activity, sterol methyl transferase I enzyme activity, sterol C4-demethylase enzyme activity, obtusifolioside C14 α -demethylase enzyme activity, sterol C5-desaturase enzyme activity, or sterol methyl transferase II enzyme activity. Thus, a plant or cell that contains introduced polynucleotide sequences other than those above, would still be considered "non-transformed."

As used herein, "peptide" and "protein" are used interchangeably and mean a compound that consists of two or more amino acids that are linked by means of peptide bonds.

I. Plant Steroid Biosynthesis

To aid the reader in understanding the present invention, descriptions of the sterol compound biosynthetic pathway are presented below. These descriptions identify enzymes useful in achieving the modifications to the biosynthesis and accumulation of sterol compounds described herein, and identify sources of nucleic acid sequences encoding these enzymes.

Various steps in the steroid compound biosynthetic pathway in plants are shown in Scheme 1, below. The numbers over the arrows refer to plant sterol compound

biosynthetic pathway enzymes and genes as indicated in Table 1.

SCHEME 1




Table 1

Plant Sterol Compound Pathway Enzymes and Genes

Enzyme	Step in Pathway	GenBank Gene ID
Acetoacetyl-CoA thiolase	1	X78116
HMG-CoA synthase	2	X83882
HMG-CoA reductase	3	X15032
		L19262
Mevalonate kinase	4	X77793
Phosphomevalonate kinase	5	Not available
Mevalonate pyrophosphate decarboxylase	6	Y14325
Isopentenyl diphosphate isomerase	7	U49259
		U47324
Farnesyl pyrophosphate synthase	8	X75789
Squalene synthase	9	AF0045 60
Squalene epoxidase	10	AB0168 83
Squalene cyclase	11	U87266
Sterol C-24 methyltransferase	12, 18	U71400
Sterol C-4 demethylase	13, 19	Not available
Cycloeucalenol- obtusifoliol isomerase	14	Not available
Sterol C-14 demethylase	15	U74319

T00290" 22/5/95

Sterol C-14 reductase	16	PCT WO 97/48793
Sterol C-8 isomerase	17	AF0303 57
Sterol C-5 desaturase	20	X90454
Sterol C-7 reductase	21	U49398
Sterol C-24 isomerase	22	Klahre et al. (1998) <i>Plant Cell</i> 10: 1677- 1690
Sterol C-24 reductase	23	Same as 22
Sterol C-22 desaturase	24	Not available
Sterol C-5 reductase	25	WO 00/61771

The plant sterol compound biosynthesis pathway has two distinct components. The early pathway reactions, leading from acetyl-CoA to squalene via mevalonic acid, are common to other isoprenoids. The later pathway reactions, leading from squalene to the major plant sterol compounds such as sitosterol, campesterol and stigmasterol, are committed biosynthetic reactions.

The early pathway reactions have been studied in
10 fungi and plants (Lees et al., *Biochemistry and
Function of Sterols*, Nes and Parish, Eds., CRC Press,
85-99 (1997); Newman and Chappell, *Biochemistry and
Function of Sterols*, Nes and Parish, Eds., CRC Press,
123-134 (1997); Bach et al., *Biochemistry and Function*

of Sterols, Nes and Parish, Eds., CRC Press, 135-150 (1997)).

Acetoacetyl CoA thiolase (EC 2.3.1.9) catalyzes the first reported reaction, which consists of the formation of acetoacetyl CoA from two molecules of acetyl CoA (Dixon et al., *J. Steroid Biochem. Mol. Biol.* 62: 165-171 (1997)). This enzyme has been purified from radish. A radish cDNA has been isolated by functional complementation in *Saccharomyces cerevisiae* (GeneBank Accession # X78116). A radish cDNA has also been screened against a cDNA library of *Arabidopsis thaliana* (Vollack and Bach, *Plant Physiology* 111: 1097-1107 (1996)).

HMGCoA synthase (EC 4.1.3.5) catalyzes the production of HMGCoA. This reaction condenses acetyl CoA with acetoacetyl CoA to yield HMGCoA. HMGCoA synthase has been purified from yeast. A plant HMGCoA synthase cDNA has also been isolated from *Arabidopsis thaliana* (Montamat et al., *Gene* 167: 197-201 (1995)).

HMGCoA reductase, also referred to as 3-hydroxy-3-methylglutaryl-coenzyme A (EC 1.1.1.34), catalyzes the reductive conversion of HMGCoA to mevalonic acid (MVA). This reaction is reported to play a role in controlling plant isoprenoid biosynthesis (Gray, *Adv. Bot. Res.* 14: 25-91 (1987); Bach et al., *Lipids* 26: 637-648 (1991); Stermer et al., *J. Lipid Res.* 35: 1133-1140 (1994). Plant HMGCoA reductase genes are often encoded by multigene families. The number of genes comprising each multigene family varies, depending on the species, ranging from two in *Arabidopsis thaliana* to at least seven in potato. Overexpression of plant HMGCoA reductase genes in transgenic tobacco plants has been reported to result in the overproduction of

09885723.062004
T00290" E2Z58860

phytosterols (Schaller et al., *Plant Physiol.* 109: 761-770 (1995)).

Mevalonate kinase (EC 2.7.1.36) catalyzes the phosphorylation of mevalonate to produce mevalonate 5-phosphate. It has been reported that mevalonate kinase plays a role in the control of isoprenoid biosynthesis (Lalitha et al., *Indian. J. Biochem. Biophys.* 23: 249-253 (1986)). A mevalonate kinase gene from *Arabidopsis thaliana* has been cloned (GeneBank accession number X77793; Riou et al., *Gene* 148: 293-297 (1994)).

Phosphomevalonate kinase (EC 2.7.4.2) (MVAP kinase) is an enzyme associated with isoprene and ergosterol biosynthesis that converts mevalonate-5-phosphate to mevalonate-5-pyrophosphate utilizing ATP (Tsay et al., *Mol. Cell. Biol.* 11: 620-631 (1991)).

Mevalonate pyrophosphate decarboxylase ("MVAPP decarboxylase") (EC 4.1.1.33) catalyzes the conversion of mevalonate pyrophosphate to isopentenyl diphosphate ("IPP"). The reaction is reported to be a decarboxylation/dehydration reaction which hydrolyzes ATP and requires Mg^{2+} . A cDNA encoding *Arabidopsis thaliana* MVAPP decarboxylase has been isolated (Toth et al., *J. Biol. Chem.* 271: 7895-7898 (1996)). An isolated *Arabidopsis thaliana* MVAPP decarboxylase gene was reported to be able to complement the yeast MVAPP decarboxylase.

Isopentenyl diphosphate isomerase ("IPP:DMAPP") (EC 5.3.3.2) catalyzes the formation of dimethylallyl pyrophosphate (DMAPP) from isopentenyl pyrophosphate (IPP). Plant IPP:DMAPP isomerase gene sequences have been reported for this enzyme. It has also been reported that IPP:DMAPP isomerase is involved in rubber

biosynthesis in a latex extract from *Hevea* (Tangpakdee
et al., *Phytochemistry* 45: 261-267 (1997)).

Farnesyl pyrophosphate synthase (EC 2.5.1.1) is a
prenyltransferase which has been reported to play a
5 role in providing polyisoprenoids for sterol compound
biosynthesis as well as a number of other pathways (Li
et al., *Gene* 17: 193-196 (1996)). Farnesyl
pyrophosphate synthase combines DMAPP with IPP to yield
geranyl pyrophosphate ("GPP"). The same enzyme
10 condenses GPP with a second molecule of IPP to produce
farnesyl pyrophosphate ("FPP"). FPP is a molecule that
can proceed down the pathway to sterol compound
synthesis, or that can be shuttled through other
pathways leading to the synthesis of quinones or
15 sesquiterpenes.

Squalene synthase (EC 2.5.1.21) reductively
condenses two molecules of FPP in the presence of Mg^{2+}
and NADPH to form squalene. The reaction involves a
head-to-head condensation, and forms a stable
20 intermediate, presqualene diphosphate. The enzyme is
subject to sterol demand regulation similar to that of
HMGCoA reductase. The activity of squalene synthase
has been reported to have a regulatory effect on the
incorporation of FPP into sterol and other compounds
25 for which it serves as a precursor (Devarenne *et al.*,
Arch. Biochem. Biophys. 349: 205-215 (1998)).

Squalene epoxidase (EC 1.14.99.7) (also called
squalene monooxygenase) catalyzes the conversion of
squalene to squalene epoxide (2,3-oxidosqualene), a
30 precursor to the initial sterol molecule in the sterol
compound biosynthetic pathway, cycloartenol. This is
the first reported step in the pathway where oxygen is
required for activity. The formation of squalene

098572-062004
"00290" E225860

epoxide is also the last common reported step in sterol biosynthesis of animals, fungi, and plants.

The later pathway of sterol compound biosynthetic steps starts with the cyclization of squalene epoxide and ends with the formation of 5 -24-alkyl sterols in plants.

2,3-oxidosqualene cycloartenol cyclase (EC 5.4.99.8) (also called cycloartenol synthase) is the first step in the sterol compound pathway that is plant-specific. The cyclization of 2,3-oxidosqualene leads to lanosterol in animals and fungi, while in plants the product is cycloartenol. Cycloartenol contains a 9,19-cyclopropyl ring. The cyclization is reported to proceed from the epoxy end in a chair-boat-chair-boat sequence that is mediated by a transient C-20 carbocationic intermediate.

S-adenosyl-L-methionine:sterol C-24 methyl transferase ("SMT1") (EC 2.1.1.41) catalyzes the transfer of a methyl group from S-adenosyl-L-methionine to the C-24 center of the sterol side chain (Nes et al. (1991) *J. Biol. Chem.* 266(23):15202-15212). This is the first of two methyl transfer reactions that have been reported to be an obligatory and rate-limiting step of the sterol compound-producing pathway in plants. The second methyl transfer reaction occurs later in the pathway after the Δ^{8-7} isomerase. The enzyme responsible for the second methyl transfer reaction is named SMTII (Bouvier-Nave, P. et al., (1997) *Eur. J. Biochem.*, 246: 518-529). An isoform, SMTII, catalyzes the conversion of cycloartenol to a $\Delta^{23(24)}$ -24-alkyl sterol, cyclosadol (Guo et al. (1996) *Tetrahed. Lett.* 37(38):6823-6826).

Sterol C-4 demethylase catalyzes the first of several demethylation reactions, which results in the removal of the two methyl groups at C-4. While in animals and fungi the removal of the two C-4 methyl groups occurs consecutively, in plants it has been reported that there are other steps between the first and second C-4 demethylations. The C-4 demethylation is catalyzed by a complex of microsomal enzymes consisting of a monooxygenase, an NAD⁺-dependent sterol 4-decarboxylase, and an NADPH-dependent 3-ketosteroid reductase.

Cycloeucalenol-obtusifoliol isomerase ("COI") catalyzes the opening of the cyclopropyl ring at C-9. The opening of the cyclopropyl ring at C-9 creates a double bond at C-8.

Sterol C-14 demethylase catalyzes demethylation at C-14, which removes the methyl group at C-14 and creates a double bond at that position. In both fungi and animals, this is the first step in the sterol synthesis pathway. Sterol 14-demethylation is mediated by a cytochrome P-450 complex.

Sterol C-14 reductase catalyzes a C-14 demethylation that results in the formation of a double bond at C-14 (Ellis et al., *Gen. Microbiol.* 137: 2627-2630 (1991)). This double bond is removed by a Δ^{14} reductase. The normal substrate is 4-methyl-8,14,24 (24¹)-trien-3 β -ol. NADPH is the normal reductant.

Sterol C-8 isomerase catalyzes a reaction that involves further modification of the tetracyclic rings or the side chain (Duratti et al., *Biochem. Pharmacol.* 34: 2765-2777 (1985)). The kinetics of the sterol isomerase-catalyzed reaction favor a $\Delta^8 \rightarrow \Delta^7$ isomerase reaction that produces a Δ^7 group.

1 Sterol C-5 desaturase catalyzes the insertion of
the Δ^5 -double bond that normally occurs at the Δ^7 -sterol
level, thereby forming a $\Delta^{5,7}$ -sterol (Parks et al.,
Lipids 30: 227-230 (1995)). The reaction has been
5 reported to involve the stereospecific removal of the
5 α and 6 α hydrogen atoms, biosynthetically derived from
the 4 pro-R and 5 pro-S hydrogens of the (+) and (-) R-
mevalonic acid, respectively. The reaction is
obligatorily aerobic, and requires NADPH or NADH. The
10 desaturase has been reported to be a multienzyme
complex present in microsomes. It consists of the
desaturase itself, cytochrome b₅, and a pyridine
nucleotide-dependent flavoprotein. The Δ^5 -desaturase is
reported to be a mono-oxygenase that utilizes electrons
15 derived from a reduced pyridine nucleotide via
cytochrome b₅.

Sterol C-7 reductase catalyzes the reduction of a
 Δ^7 -double bond in $\Delta^{5,7}$ -sterols to generate the
corresponding Δ^5 -sterol. It has been reported that the
20 mechanism involves, like many other sterol enzymes, the
formation of a carbocationic intermediate via
electrophilic "attack" by a proton.

Sterol C-24(28) isomerase catalyzes the reduction
of a $\Delta^{24(28)}$ - Δ^{24} , a conversion that modifies the side
25 chain. The product is a $\Delta^{24(25)}$ -24-alkyl sterol. Sterol
C-24 reductase catalyzes the reduction of the ²⁴⁽²⁵⁾
double bond at C-24, which produces sitosterol.
Recently, Klahre et al. ((1998) *Plant Cell* 10:1677-
1690) discovered that both the isomerization and
30 reduction steps are catalyzed by an enzyme coded by the
same gene, i.e., *DIM/DWF1*.

sterol C-22 desaturase (EC 2.7.3.9) catalyzes the formation of a double bond at C-22 on the side chain. This formation of a double bond at C-22 on the side chain marks the end of the sterol compound biosynthetic pathway, and results in the formation of stigmasterol (Benveniste (1986) *Annu. Rev. Plant Physiol.* 37:275-308). The C-22 desaturase in yeast, which is the reported final step in the biosynthesis of ergosterol in that organism, requires NADPH and molecular oxygen. In addition, the reaction is also reported to involve a cytochrome P450 that is distinct from a cytochrome P450 participating in demethylation reactions (Lees et al. (1995) *Lipids* 30: 221-226).

Phytosterols are biogenetic precursors of brassinosteroids, steroid alkaloids, steroid sapogenins, ecdysteroids, and steroid hormones. This precursor role of phytosterols is often described as a "metabolic" function. A common transformation of free sterols in tissues of vascular plants is the conjugation at the 3-hydroxy group of sterols with long-chain fatty acids to form steryl esters, or with a sugar, usually with a single molecule of β -D-glucose, to form steryl glycosides. Some of the steryl glycosides are additionally esterified, at the 6-hydroxy group of the sugar moiety, with long-chain fatty acids to form acylated steryl glycosides.

The existence of several enzymes that are specifically associated with the synthesis and breakdown of conjugated sterols has been reported (Wojciechowski, *Physiology and Biochemistry of Sterols*, eds. Patterson, Nes, AOCs Press, 361 (1991)). Enzymes involved in this process include: UDPGlc:Sterol glucosyltransferase, phospho(galacto)glyceride steryl

T00290" E2253360

glucoside acyltransferase, and sterylglycoside and
sterylester hydrolases.

UDPGlc:sterol glucosyltransferase (EC 2.4.1.173)
catalyzes glucosylation of phytosterols by glucose
5 transfer from UDP-glucose ("UDPGl"). The formation of
steryl glycosides can be measured using UDP- [¹⁴C]glucose
as the substrate. Despite certain differences in their
specificity patterns, all reported UDPGlc:sterol
glucosyltransferases preferentially glucosylate only
10 sterols or sterol-like molecules that contain a C-3
hydroxy group, a β -configuration, and which exhibit a
planar ring. It has been reported that UDPGlc:sterol
glucosyltransferases are localized in the microsomes.

Phospho(galacto)glyceride steryl glucoside
15 acyltransferase catalyzes the formation of acylated
steryl glycosides from the substrate steryl glycoside
by transfer of acyl groups from some membranous polar
acyllipids to steryl glycoside molecules.

Acylglycerol:sterol acyltransferase (EC 2.3.1.26)
20 catalyzes the reaction wherein certain acylglycerols
act as acyl donors in a phytosterol esterification. In
plants, the activity of acylglycerol:sterol
acyltransferase is reported to be associated with
membranous fractions. A pronounced specificity for
25 shorter chain unsaturated fatty acids was reported for
all acyltransferase preparations studied in plants.
For example, acylglycerol:sterol acyltransferases from
spinach leaves and mustard roots can esterify a number
of phytosterols.

30 Sterylglycoside and sterylester hydrolases ("SG-
hydrolases") catalyze the enzymatic hydrolysis of
sterylglycosides to form free sterols. The SG-
hydrolase activity is not found in mature, ungerminated

0988723-062004

5 Enzymatic hydrolysis of sterylesters in germinating
seeds of mustard, barley and corn is reported to be low
in dormant seeds, but increases during the first ten
days of germination. This activity is consistent with a
decrease in sterylesters and an increase in free
10 sterols over the same temporal period.

In order to obtain seed producing oil containing elevated levels of phytosterols and phytosterol esters such as sitosterol and sitosterol esters, these recombinant constructs or expression cassettes can be introduced into plant cells by any number of conventional means known in the art and regenerated into fertile transgenic plants. The genome of such plants can then comprise introduced DNA encoding various steroid pathway enzymes, alone or in combination, that achieves the desirable effect of enhancing the levels of phytosterols, phytosterol esters, mixtures thereof in the oil of seed thereof.

Preferably, the genome can comprise introduced DNA encoding a HMG CoA reductase enzyme and an introduced DNA encoding one or more of a squalene epoxidase, a sterol methyl transferase I, a sterol C4-demethylase, an obtusifoliol C14 α -demethylase, a sterol C5-desaturase, a sterol methyl transferase II. In each case, the foregoing introduced DNAs can be operatively

linked to regulatory signals that cause seed-specific expression thereof.

The present invention encompasses not only such transgenic plants, but also transformed plant cells, including cells and seed of such plants, as well as progeny of such plants, for example produced from the seed. Transformed plant cells and cells of the transgenic plants encompassed herein can be grown in culture for a time and under appropriate conditions to produce oil containing elevated levels of phytosterols and/or phytostanols and their corresponding esters. Alternatively, the phytosterols, phytostanols, and their corresponding esters can be isolated directly from the cultures.

In addition, of course, seed obtained from the transgenic, progeny, hybrid, etc., plants disclosed herein can be used in methods for obtaining oil containing phytosterols, phytosterol esters, phytostanols, phytostanol esters, or mixtures thereof employing extraction and processing procedures known in the art. Note, in this regard, Kochhar (1983) *Prog. Lipid Res.* 22: 161-188.

The present invention also encompasses a method of producing a plant that accumulates an elevated level of sitosterol, at least one sitosterol ester, sitostanol, at least one sitostanol ester, or mixtures thereof, in seeds thereof compared to seeds of a corresponding plant comprising no introduced DNA encoding a polypeptide or protein that affects the biosynthesis of sterols, phytosterols, phytosterol esters, phytostanols, phytostanol esters, or combinations thereof, comprising sexually crossing a transgenic plant of the present invention with such a corresponding plant. The latter can be a non-

Variable	Mean	SD	Min	Max
Age	38.5	12.5	18	65
Gender	0.5	0.5	0	1
Marital status	0.7	0.5	0	1
Education	12.5	2.5	9	16
Income	1500	500	500	3000
Health status	0.8	0.4	0	1
Exercise frequency	2.5	1.5	0	5
Stress level	3.5	1.5	1	5
Sleep quality	4.0	1.0	2	5
Work satisfaction	3.0	1.0	1	5
Life satisfaction	4.5	1.0	2	5

Variable	Mean	SD	Min	Max
Age	38.5	12.5	18	65
Gender	0.5	0.5	0	1
Marital status	0.7	0.5	0	1
Education	12.5	2.5	9	16
Income	1500	500	500	3000
Health status	0.8	0.4	0	1
Exercise frequency	2.5	1.5	0	5
Stress level	3.5	1.5	1	5
Sleep quality	4.0	1.0	2	5
Work satisfaction	3.0	1.0	1	5
Life satisfaction	4.5	1.0	2	5

A process of increasing the formation of steroid pathway products in a transformed host cell as compared to an otherwise identical non-transformed host cell comprising the following steps. A host cell is transformed with a recombinant vector comprising (a) as operably linked components in the 5' to 3' direction, a promoter, a DNA sequence encoding a polypeptide exhibiting 3-hydroxy-3-methylglutaryl-Coenzyme A reductase enzyme activity, and a transcription termination signal sequence; and (b) as operably linked components in the 5' to 3' direction, a promoter, a DNA sequence encoding a steroid pathway enzyme, and a transcription termination signal sequence. The steroid pathway enzyme is a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifolioside C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, and a sterol methyl

transferase II enzyme. The transformed plant cell is regenerated into a transgenic plant.

A plant contemplated by this invention is a vascular, multicellular higher plant. Such higher plants will hereinafter be usually referred to simply as "plants". Such "plants" include both complete entities having leaves, stems, seeds, roots and the like as well as callus and cell cultures that are monocotyledonous and dicotyledonous. Dicotyledonous plants are a preferred embodiment of the present invention.

Preferred plants are members of the Solanaceae, Leguminosae, Ammiaceae, Brassicaceae, Gramineae, Carduaceae and Malvaceae families. Exemplary plant members of those families are tobacco, petunia and tomato (Solanaceae), soybean and alfalfa (Leguminosae), carrot (Ammiaceae), corn, maize and barley (Gramineae), *Arabidopsis* (Brassicaceae), guayule (Carduaceae), and cotton (Malvaceae). A preferred plant is tobacco of the strain *Nicotiana tabacum* (*N. Tabacum*), cotton of the strain Coker line 312-5A, soybean of the strain *Glycine max*, alfalfa of the strain RYSI or tomato of the strain *Lycopersicon esculentum*. Other plants include canola, maize and rape.

A transgenic plant contemplated by this invention is produced by transforming a plant cell or protoplast with an added, exogenous structural gene that encodes a polypeptide having HMG-CoA reductase activity and an exogenous structural gene that encodes at least one polypeptide having steroid pathway enzyme activity to produce a transformed plant cell, and regenerating a transgenic plant from the transformed plant cell. The encoded polypeptide is expressed both in the transformed plant cell or protoplast and the resulting

transgenic plant. (The phrase "plant cell" will hereinafter be used to include a plant protoplast, except where plant protoplasts are specifically discussed).

5 A non-transgenic plant that serves as the source of the plant cell that is transformed, i.e. the precursor cell, is referred to herein as a "native, non-transgenic" plant. The native, non-transgenic plant is of the same strain as the formed transgenic
10 plant.

Sterol production in a transgenic plant of the present invention is increased by increasing the activity of the enzyme HMG-CoA reductase, which enzyme catalyzes the conversion of 3-hydroxy-3-methylglutaryl
15 Coenzyme A (HMG-CoA) to mevalonate and the activity of at least one other steroid pathway enzyme. As used herein, the term "specific activity" means the activity normalized to cellular protein content.

HMG-CoA reductase activity is increased by
20 increasing the amount (copy number) of a gene encoding a polypeptide having HMG-CoA reductase catalytic activity. Expression of the increased amount of that encoded structural gene enhances the activity of that enzyme.

25 The amount of the expressed gene is increased by transforming a plant cell with a recombinant DNA molecule comprising a vector operatively linked to a DNA segment that encodes a polypeptide having HMG-CoA reductase activity, and a promoter suitable for
30 driving the expression of that polypeptide in that plant cell, and culturing the transformed plant cell into a transgenic plant. Such a polypeptide includes intact as well as a catalytically active, truncated HMG-CoA reductase proteins.

09885723-062004
T00290" E2Z58860

Thus, a transformed plant cell and a transgenic plant have one or more added, exogenous genes that encode a polypeptide having HMG-CoA reductase activity and at least one other steroid pathway enzyme activity relative to a native, non-transgenic plant or untransformed plant cell of the same type. As such, a transformed plant cell or transgenic plant can be distinguished from an untransformed plant cell or native, nontransgenic plant by standard technology such as agarose separation of DNA fragments or mRNAs followed by transfer and appropriate blotting with DNA or RNA, e.g., Southern or Northern blotting, or by use of polymerase chain reaction technology, as are well known. Relative HMG-CoA reductase activity of the transformed cell or transgenic plant with untransformed cells and native, non-transgenic plants or cell cultures therefrom can also be compared, with a relative activity for that enzyme of about 1.5:1 for transgenic (transformed) to native (untransformed) showing transformation. Higher relative activity ratios such as about 15:1 have also been observed.

Sterol accumulation can also be used to distinguish between native, non-transgenic and transgenic plants. A transgenic plant has at least about twice the total sterol content as a native, non-transgenic plant where a single added gene is present. Greater differences up to about forty-fold have also been observed.

Sitostanol, sitostanol ester, and tocopherol biosynthesis and accumulation in plants can be modified in accordance with the present invention by variously expressing the nucleic acid coding sequences discussed above, alone or in combination, as described herein. The expression of sequences encoding sterol

methytransferase II enzymes facilitates the production of plants in which the biosynthesis and accumulation of campesterol, campestanol, and their esters can be reduced as these enzymes shunt sterol intermediates
5 away from campesterol, and toward sitosterol and sitostanol.

III. DNA Encoding Useful Polypeptides

The present invention contemplates a recombinant
10 construct or a recombinant vector that contains a DNA sequence encoding a polypeptide exhibiting 3-hydroxy-3-methylglutaryl-Coenzyme A (HMG-CoA) reductase activity and a DNA sequence encoding a polypeptide exhibiting the activity of a steroid pathway enzyme. Each
15 polypeptide-encoding DNA sequence is operably linked in the 5' to 3' direction independent of the other sequence. Each DNA sequence in the 5' to 3' direction comprises a promoter, then the DNA sequence encoding the polypeptide then a transcription termination signal
20 sequence. The steroid pathway enzyme is a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifoliol C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, or a sterol methyl transferase II enzyme. It is
25 contemplated that HMG-CoA reductase and steroid pathway enzyme activities come from a mutant or truncated form of those enzymes, such as a truncated HMG-CoA reductase lacking the transmembrane region while retaining a functional catalytic domain. Several HMG CoA reductase
30 sequences are known in the art. An amino acid alignment for these is shown in FIG. 32. The sources of the sequences used in building the multiple alignment are listed in Table 5.

036672-062004
T00290"E245850

Table 5. Sources of Sequences Used In Building
The Multiple Alignment

methanobac	swissprot:hmdh_meth	Begin:1	End:397	O26662 methanobacterium thermoautotrophicur
methanococ	swissprot:hmdh_metja	Begin:1	End:405	Q58116 methanococcus jannaschii. 3-hydroxy-3
halobacter	swissprot:hmdh_halvo	Begin:1	End:403	Q59468 halobacterium volcanii (haloferax volcar
sulfolobus	swissprot:hmdh_sulso	Begin:1	End:409	O08424 sulfolobus solfataricus. 3-hydroxy-3-me
yeast2	gp_pln1:yschmgcr2_1	Begin:1	End:1045	M22255 Saccharomyces cerevisiae Yeast HMG
yeast1	gp_pln1:yschmgcr1_1	Begin:1	End:1054	M22002 Saccharomyces cerevisiae Yeast HMG
phycomyces	swissprot:hmdh_phybl	Begin:1	End:105	Q12649 phycomyces blakesleeanus. 3-hydroxy-
fusarium	swissprot:hmdh_fusmo	Begin:1	End:976	Q12577 fusarium moniliforme (gibberella fujikurc
candida	gp_pln1:ab012603_1	Begin:1	End:934	AB012603 Candida utilis Candida utilis HMG mF
dictyoste2	swissprot:hmd2_dicdi	Begin:1	End:481	P34136 dictyostelium discoideum (slime mold). :
wheat1	pir2:pq0761	Begin:1	End:150	hydroxymethylglutaryl-CoA reductase (NADPH)
rice	swissprot:hmdh_orysa	Begin:1	End:509	P48019 oryza sativa (rice). 3-hydroxy-3-methylg
corn	sp_plant:o24594	Begin:1	End:579	O24594 zea mays (maize). 3-hydroxy-3-methylg
wheat3	pir2:pq0763	Begin:1	End:150	hydroxymethylglutaryl-CoA reductase (NADPH)
wheat2	pir2:pq0762	Begin:1	End:150	hydroxymethylglutaryl-CoA reductase (NADPH)
soybean	gmtx6:30820_1r59f1	Begin:101	End:259	from proprietary soy sequence database
rubbertre3	swissprot:hmd3_hevbr	Begin:1	End:586	Q00583 hevea brasiliensis (para rubber tree). 3-
rosyperiwi	swissprot:hmdh_catro	Begin:1	End:601	Q03163 catharanthus roseus (rosy periwinkle) (r
tomato	swissprot:hmd2_lyces	Begin:1	End:602	P48022 lycopersicon esculentum (tomato). 3-hy
woodtobacc	swissprot:hmdh_nicsy	Begin:1	End:604	Q01559 nicotiana sylvestris (wood tobacco). 3-h
potato	gp_pln1:potmgri_1	Begin:1	End:596	L01400 Solanum tuberosum Potato hydroxymeth
radish	sp_plant:q43826	Begin:1	End:573	Q43826 raphanus sativus (radish). hydroxymeth
arabidopsis1	gp_pln1:athhmgcoar_1	Begin:1	End:592	L19261 Arabidopsis thaliana Arabidopsis thalian
cucumismel	gp_pln1:ab021862_1	Begin:1	End:587	AB021862 Cucumis melo Cucumis melo mRNA
rubbertre2	swissprot:hmd2_hevbr	Begin:1	End:210	P29058 hevea brasiliensis (para rubber tree). 3-
rubbertre1	swissprot:hmd1_hevbr	Begin:1	End:575	P29057 hevea brasiliensis (para rubber tree). 3-
camptothec	swissprot:hmdh_camac	Begin:1	End:593	P48021 camptotheca acuminata. 3-hydroxy-3-m
arabidops2	swissprot:hmd2_arath	Begin:1	End:562	P43256 arabidopsis thaliana (mouse-ear cress).
chineseham	swissprot:hmdh_crigr	Begin:1	End:887	P00347 cricetus griseus (chinese hamster). 3-
chineseha2	gp_rod:cruhmg14_1	Begin:1	End:887	L00183 Cricetus sp. Hamster 3-hydroxy-3-met
syrianhamst	gp_rod:hamhmgcob_1	Begin:1	End:887	M12705 Mesocricetus auratus Syrian hamster 3
rat	swissprot:hmdh_rat	Begin:1	End:887	P51639 rattus norvegicus (rat). 3-hydroxy-3-met
rabbit	swissprot:hmdh_rabit	Begin:1	End:888	Q29512 oryctolagus cuniculus (rabbit). 3-hydrox
human	gp_pri2:humhmgcoa_1	Begin:1	End:888	M11058 Homo sapiens Human 3-hydroxy-3-met
mouse	gp_rod:mushmgcoa_1	Begin:1	End:224	M62766 Mus musculus Mouse HMG-CoA reduc
xenopus	swissprot:hmdh_xenla	Begin:1	End:883	P20715 xenopus laevis (african clawed frog). 3-
seaurchin	swissprot:hmdh_strpu	Begin:1	End:932	P16393 stronglylacentrotus purpuratus (purple s
cockroach	swissprot:hmdh_blage	Begin:1	End:856	P54960 blattella germanica (german cockroach)
drosophila	swissprot:hmdh_drome	Begin:1	End:916	P14773 drosophila melanogaster (fruit fly). 3-hy
dictyoste1	swissprot:hmd1_dicdi	Begin:1	End:552	P34135 dictyostelium discoideum (slime mold). :
schistosom	swissprot:hmdh_schma	Begin:1	End:948	P16237 schistosoma mansoni (blood fluke). 3-h
archaeoglo	swissprot:hmdh_arclu	Begin:1	End:436	O28538 archaeoglobus fulgidus. 3-hydroxy-3-mi
pseudomonas	gp_bct1:psehmgcoa_1	Begin:1	End:428	M24015 Pseudomonas mevalonii P.mevalonii HI

5

These sequences, and their truncated counterparts, are useful in the present invention. Examples of such preferred HMG CoA reductases include the truncated rubber and *Arabidopsis* HMG CoA reductases disclosed herein.

10

Other enzyme-encoding DNAs can be introduced into plants to elevate even further the levels of desirable $\Delta 5$ sterols and their reduced stanol counterparts as well as other phytosterols and tocopherols. Thus, the

DNA sequences contemplated for use in the present invention, which can be used alone or in various combinations as discussed below, include, but are not limited to, those encoding the following enzymes: 3-hydroxysteroid oxidases; steroid 5reductases; sterol methyltransferases; sterol acyltransferases; and S-adenosylmethionine-dependent α -tocopherol methyltransferases.

In each case, the sequences encoding these enzymes can comprise an expression cassette comprising, operably linked in the 5' to 3'direction, a seed-specific promoter, the enzyme coding sequence, and a transcriptional termination signal sequence functional in a plant cell such that the enzyme is successfully expressed. For use in the methods disclosed herein, the recombinant constructs or expression cassettes can be incorporated in a vector, for example a plant expression vector. Such vectors can be transformed into host cells such as bacterial cells, for example during the preparation or modification of the recombinant constructs, and plant cells. Thus, the invention encompasses plants and seeds comprising such transformed plant cells.

It will be apparent to those of skill in the art that the nucleic acid sequences set forth herein, either explicitly, as in the case of the sequences set forth above, or implicitly with respect to nucleic acid sequences generally known and not present herein, can be modified due to the built-in redundancy of the genetic code and noncritical areas of the polypeptide that are subject to modification and alteration. In this regard, the present invention contemplates allelic

[illegible]

Thus, as long as the DNA segment is free of such interfering DNA sequences, a DNA segment of the invention can be up to 15,000 base pairs in length. The maximum size of a recombinant DNA molecule, particularly a plant integrating vector, is governed mostly by convenience and the vector size that can be accommodated by a host cell, once all of the minimal DNA sequences required for replication and expression, when desired, are present. Minimal vector sizes are well known.

Also encompassed by the present invention are
30 nucleotide sequences biologically functionally
equivalent to those disclosed herein, that encode
conservative amino acid changes within the amino acid
sequences of the presently disclosed enzymes, producing
"silent" changes therein. Such nucleotide sequences

contain corresponding base substitutions based upon the genetic code compared to the nucleotide sequences encoding the presently disclosed enzymes. Substitutes for an amino acid within the enzyme sequences disclosed
5 herein is selected from other members of the class to which the naturally occurring amino acid belongs. Amino acids can be divided into the following four groups: (1) acidic amino acids; (2) basic amino acids; (3) neutral polar amino acids; and (4) neutral non-
10 polar amino acids. Representative amino acids within these various groups include, but are not limited to: (1) acidic (negatively charged) amino acids such as aspartic acid and glutamic acid; (2) basic (positively charged) amino acids such as arginine, histidine, and
15 lysine; (3) neutral polar amino acids such as glycine, serine, threonine, cysteine, cystine, tyrosine, asparagine, and glutamine; and (4) neutral nonpolar (hydrophobic) amino acids such as alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan,
20 and methionine.

A. HMG-CoA Reductase

The introduction of an HMG CoA reductase gene into a cell results in a higher carbon throughput through
25 the steroid synthesis pathway. The introduction of a truncated HMG CoA reductase gene (lacking the transmembrane region, resulting in a soluble HMG CoA reductase enzyme) provides higher HMG CoA reductase activity and thus increased delta-5 steroid compound
30 production over the same case with an introduced full-length HMG CoA reductase gene. A useful truncated HMG CoA reductase nucleic acid encodes at least the catalytic domain.

[illegible]

The catalytic region contains the active site of the HMG-CoA reductase enzyme and comprises about forty percent of the COOH-terminal portion of intact HMG-CoA reductase enzyme.

25 The linker region connects the catalytic and
 membrane-binding regions, and constitutes the remaining
 about ten percent of the intact enzyme.

As discussed in greater detail below, only the catalytic region of HMG-CoA reductase is needed herein to provide the desired enzyme activity. Thus, an exogenous structural gene that encodes a polypeptide corresponding to that catalytic region is the minimal HMG Co A reductase gene required for transforming plant cells in addition to one of the steroid pathway enzymes

discussed below. The present invention therefore contemplates use of both intact and truncated structural genes that encode a polypeptide having HMG-CoA reductase activity.

5 A structural gene encoding a polypeptide having HMG-CoA reductase activity can be obtained or constructed from a variety of sources and by a variety of methodologies. See, e.g. Carlson et al., *Cell*, 28:145 (1982); Rine et al., *Proc. Natl. Acad. Sci. USA*, 10 80:6750 (1983). Exemplary of such structural genes are the mammalian and yeast genes encoding HMG-CoA reductase or the catalytic region thereof.

The disclosures of Chappell, et al., U.S. Patent No. 5,349,126, are incorporated in full herein by 15 reference. The mammalian genome contains a single gene encoding HMG-CoA reductase. The nucleotide sequence of the hamster and human gene for HMG-CoA reductase have been described in Chappell et al. A composite nucleotide sequence of DNA corresponds to the mRNA SEQ 20 ID NO:1 of Chappell et al., as well as the derived amino acid residue sequence SEQ ID NO:2 of Chappell et al., for hamster HMG-CoA reductase is provided in Fig. 2 of Chappell et al, reprinted from Chin et al., *Nature*, 308:613 (1984). The composite nucleotide 25 sequence of Fig. 2, SEQ ID NO:1 of Chappell et al., comprising about 4768 base pairs, includes the nucleotide sequence encoding the intact hamster HMG-CoA reductase enzyme.

Intact hamster HMG-CoA reductase comprises about 30 887 amino acid residues (SEQ ID NO:2 of Chappell et al.). A structural gene encoding an intact hamster HMG-CoA reductase enzyme of 887 amino acid residues comprises base pairs from about nucleotide position 164

098837-062001
F00290" E3/58860

to about nucleotide position 2824 of SEQ ID NO:1 of Chappell et al.

A preferred structural gene is one that encodes a polypeptide corresponding to only the catalytic region of the enzyme. Two catalytically active segments of hamster HMG-CoA reductase have been defined. Liscum et al., *J. Biol. Chem.*, 260(1):522 (1985). One segment containing a catalytic region has an apparent molecular weight of 62 kDa and comprises amino acid residues from about position 373 to about position 887. A second segment containing a catalytic region has an apparent molecular weight of 53 kDa segment and comprises amino acid residues from about position 460 to about position 887. The 62 kDa catalytically active segment is encoded by base pairs from about nucleotide position 1280 to about nucleotide position 2824 of SEQ ID NO:1 of Chappell et al. The 53 kDa catalytically active segment is encoded by base pairs from about nucleotide position 1541 to about nucleotide position 2824 of SEQ ID NO:1 of Chappell et al.

In a preferred embodiment, the utilized structural gene encodes the catalytic region and at least a portion of the linker region of HMG-CoA reductase. The linker region of hamster HMG-CoA reductase comprises amino acid residues from about position 340 to about position 373 or from about position 340 to about position 460, depending upon how the catalytic region is defined. These linker regions are encoded by base pairs from about nucleotide position 1180 to about nucleotide position 1283 or from about position 1180 to about position 1540, respectively of SEQ ID NO:1 of Chappell et al. The structural gene encoding the linker region is operatively linked to the structural gene encoding the catalytic region.

In one particularly preferred embodiment, a structural gene encoding a catalytically active, truncated HMG-CoA reductase enzyme can optionally contain base pairs encoding a small portion of the membrane region of the enzyme.

A structural gene encoding a polypeptide comprising a catalytically active, truncated or intact HMG-CoA reductase enzyme from other organisms such as yeast can also be used in accordance with the present invention.

Yeast cells contain two genes encoding HMG-CoA reductase. The two yeast genes, designated HMG1 and HMG2, encode two distinct forms of HMG-CoA reductase, designated HMG-CoA reductase 1 SEQ ID NO:3 of Chappell et al. are presented in Fig. 3 of Chappell et al., are taken from Basson et al. *Mol. Cell Biol.*, 8(9):3797 (1988). The nucleotide base sequences of HMG2 SEQ ID NO:5 of Chappell et al. as well as the amino acid residue sequence of HMG-CoA reductase 2 SEQ ID NO:6 of Chappell et al. are set forth therein in the Sequence Listing.

The entire HMG1 gene comprises about 3360 base pairs SEQ ID NO:3 of Chappell et al. Intact HMG-CoA reductase 1 comprises an amino acid sequence of about 1054 amino acid residues SEQ ID NO:4 of Chappell et al. Thus, the minimal portion of the HMG1 gene that encodes an intact enzyme comprises base pairs from about nucleotide position 121 to about position 3282 of Fig. 3, SEQ ID NO:3 of Chappell et al.

The entire HMG2 gene comprises about 3348 base pairs SEQ ID NO:5 of Chappell et al. Intact HMG-CoA reductase 2 comprises about 1045 amino acid residues SEQ ID NO:6 of Chappell et al. Thus, the minimal portion of HMG2 gene that encodes intact HMG-CoA

reductase 2 comprises base pairs from about nucleotide position 121 to about position 3255 of SEQ ID NO:5 of Chappell et al.

By analogy to the truncated hamster structural gene, structural genes encoding polypeptides comprising catalytically active, truncated HMG-CoA reductase enzymes from yeast can also be used in accordance with the present invention.

The catalytic region of HMG-CoA reductase 1 comprises amino acid residues from about residue 618 to about residue 1054: i.e., the COOH-terminus. A structural gene that encodes the catalytic region comprises base pairs from about nucleotide position 1974 to about position 3282 of Fig. 3 of Chappell et al.

The linker region of HMG-CoA reductase 1 comprises an amino acid sequence from about residue 525 to about residue 617. A structural gene that encodes the linker region comprises nucleotides from about position 1695 to about position 1973 of Fig. 3 of Chappell et al. A structural gene encoding the linker region of the enzyme operatively linked to the structural gene encoding the catalytic region of the enzyme.

Also by analogy to the truncated hamster gene, a truncated HMG1 gene can optionally contain nucleotide base pair sequences encoding a small portion of the membrane-binding region of the enzyme. Such a structural gene preferably comprises base pairs from about nucleotide position 121 to about position 147 and from about position 1695 to about position 3282 of Fig. 3 of Chappell et al.

A construct similar to those above from an analogous portion of yeast HMG-CoA reductase 2 can also be utilized.

5

10

15

15

20

25

30

Q12649 phycomyces blakesleeanus. 3-hydroxy-3-methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-coa reductase) (fragment). 11/1997

Q12577 fusarium moniliforme (gibberella
fujikuroi). 3-hydroxy-3-methylglutaryl-coenzyme a
reductase (ec 1.1.1.34) (hmg- coa reductase). 11/1997

AB012603 Candida utilis Candida utilis HMG mRNA
5 for HMG-CoA reductase, complete cds. 7/1998

P34136 dictyostelium discoideum (slime mold). 3-
hydroxy-3-methylglutaryl-coenzyme a reductase 2 (ec
1.1.1.34) (hmg- coa reductase 2) (fragment).35735

PQ0761 hydroxymethylglutaryl-CoA reductase (NADPH)
10 (EC 1.1.1.34) (HMGR 10) - wheat (fragment)

P48019 oryza sativa (rice). 3-hydroxy-3-
methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-
coa reductase) (fragment). 2/1996

O24594 zea mays (maize). 3-hydroxy-3-
15 methylglutaryl coenzyme a reductase (ec 1.1.1.88).
5/1999

PQ0763 hydroxymethylglutaryl-CoA reductase (NADPH)
(EC 1.1.1.34) (HMGR 23) - wheat (fragment)

PQ0762 hydroxymethylglutaryl-CoA reductase (NADPH)
20 (EC 1.1.1.34) (HMGR 18) - wheat (fragment)
from proprietary soy sequence database

Q00583 hevea brasiliensis (para rubber tree). 3-
hydroxy-3-methylglutaryl-coenzyme a reductase 3 (ec
1.1.1.34) (hmg- coa reductase 3). 7/1998

25 Q03163 catharanthus roseus (rosy periwinkle)
(madagascar periwinkle). 3-hydroxy-3-methylglutaryl-
coenzyme a reductase (ec 1.1.1.34) (hmg-coa reductase).
7/1998

P48022 lycopersicon esculentum (tomato). 3-
30 hydroxy-3-methylglutaryl-coenzyme a reductase 2 (ec
1.1.1.34) (hmg- coa reductase 2). 7/1998

Q01559 nicotiana glauca (wood tobacco). 3-
hydroxy-3-methylglutaryl-coenzyme a reductase (ec
1.1.1.34) (hmg-coa reductase). 7/1998

1002990 22/58850

L01400 *Solanum tuberosum* Potato
hydroxymethylglutaryl coenzyme A reductase (hmgr) mRNA,
complete cds; putative. 4/1996

Q43826 *raphanus sativus* (radish).

- 5 hydroxymethylglutaryl-coa reductase (ec 1.1.1.34)
(hydroxymethylglutaryl-coa reductase (nadph)) (3-
hydroxy-3-methylglutaryl-coenzyme a red

L19261 *Arabidopsis thaliana* *Arabidopsis thaliana*
HMG-coA reductase gene, complete cds. 4/1994

- 10 AB021862 *Cucumis melo* *Cucumis melo* mRNA for HMG-
CoA reductase, complete cds; putative. 1/1999

P29058 *hevea brasiliensis* (para rubber tree). 3-
hydroxy-3-methylglutaryl-coenzyme a reductase 2 (ec
1.1.1.34) (hmg- coa reductase 2) (fragment).35735

- 15 P29057 *hevea brasiliensis* (para rubber tree). 3-
hydroxy-3-methylglutaryl-coenzyme a reductase 1 (ec
1.1.1.34) (hmg- coa reductase 1). 7/1998

- P48021 *camptotheca acuminata*. 3-hydroxy-3-
methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-
20 coa reductase). 11/1997

P43256 *arabidopsis thaliana* (mouse-ear cress). 3-
hydroxy-3-methylglutaryl-coenzyme a reductase 2 (ec
1.1.1.34) (hmg- coa reductase 2) (hmgr2). 7/1998

- P00347 *cricetulus griseus* (chinese hamster). 3-
25 hydroxy-3-methylglutaryl-coenzyme A reductase (ec
1.1.1.34) (hmg-coA reductase). 11/1997

- L00183 *Cricetulus* sp. Hamster 3-hydroxy-3-
methylglutaryl coenzyme A (HMG CoA) reductase gene,
exons 19 and 20; 3-hydroxy-3-methylglutaryl coenzyme A
30 (HMG CoA). 4/1993

M12705 *Mesocricetus auratus* Syrian hamster 3-
hydroxy-3-methylglutaral coenzyme A reductase (HMG-CoA
reductase) mRNA, complete cds; 3-hydroxy-3-
methylglutaral coenzyme A red

0985723 062004
T00290 E245866

P51639 rattus norvegicus (rat). 3-hydroxy-3-methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-coa reductase). 12/1998

Q29512 oryctolagus cuniculus (rabbit). 3-hydroxy-
5 3-methylglutaryl-coenzyme a reductase (ec 1.1.1.34)
(hmq-coa reductase). 7/1999

M11058 Homo sapiens Human 3-hydroxy-3-methylglutaryl coenzyme A reductase mRNA, complete cds; 3-hydroxy-3-methylglutaryl coenzyme A reductase.11/1994

10 M62766 Mus musculus Mouse HMG-CoA reductase mRNA,
3' end. 4/1993

P20715 xenopus laevis (african clawed frog). 3-hydroxy-3-methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-coa reductase). 11/1997

15 P16393 stronglylocentrotus purpuratus (purple sea
urchin). 3-hydroxy-3-methylglutaryl-coenzyme a
reductase (ec 1.1.1.34) (hmg-coa reductase). 11/1997

P54960 blattella germanica (german cockroach). 3-hydroxy-3-methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-coa reductase). 11/1997

P14773 drosophila melanogaster (fruit fly). 3-hydroxy-3-methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-coa reductase). 12/1998

P34135 dictyostelium discoideum (slime mold). 3-
25 hydroxy-3-methylglutaryl-coenzyme a reductase 1 (ec
1.1.1.34) (hmg- coa reductase 1). 11/1997

P16237 schistosoma mansoni (blood fluke). 3-hydroxy-3-methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-coa reductase). 7/1998

30 O28538 archaeoglobus fulgidus. 3-hydroxy-3-
methylglutaryl-coenzyme a reductase (ec 1.1.1.34) (hmg-
coa reductase). 12/1998

[illegible]

[illegible]

5

10

i

15

20

25

30

Squalene Epoxidase, also known as squalene monooxygenase is enzyme reference number 1.14.99.7, Enzyme Nomenclature 1992, p. 146.

Several squalene epoxidase enzymes are known to the art. These include *Arabidopsis* squalene epoxidase protein sequence Accession No. AC004786 (SEQ ID NO:1), *Arabidopsis* squalene epoxidase Accession No. N64916 (SEQ ID NO:2), and *Arabidopsis* squalene epoxidase Accession No. T44667 (SEQ ID NO:3). Japanese patent application No. 07194381 A discloses a DNA encoding a mammalian squalene epoxidase.

In order to facilitate the modifications to sterol biosynthesis and accumulation described herein, the present invention also provides an isolated DNA molecule, comprising a nucleotide sequence selected from the group consisting of:

(a) *Arabidopsis* squalene epoxidase from clone ID
ATA506263 disclosure SEQ ID NO:4, clone ID ATA304243
disclosure SEQ ID NO:6, clone ID ATA102071 disclosure
20 SEQ ID NO: 8, clone ATA504158 disclosure SEQ ID NO:10,
or the complement thereof;

(b) a nucleotide sequence that hybridizes to said nucleotide sequence of (a) under a wash stringency equivalent to 0.5X SSC to 2X SSC, 0.1% SDS, at 55-65°C, and which encodes a polypeptide having squalene epoxidase enzymatic activity substantially similar to that of the disclosed squalene epoxidase;

(c) a nucleotide sequence encoding the same genetic information as said nucleotide sequence of (a), but which is degenerate in accordance with the degeneracy of the genetic code; and

(d) a nucleotide sequence encoding the same genetic information as said nucleotide sequence of (b),

but which is degenerate in accordance with the degeneracy of the genetic code.

An additional aspect of the invention is the recombinant constructs and vectors (pMON48343, Fig. 30; pMON43844, Fig. 31) comprising nucleic acid sequences encoding the novel squalene epoxidase, as well as a method of producing the novel squalene epoxidase, comprising culturing a host cell transformed with the novel constructs or vectors for a time and under conditions conducive to the production of the squalene epoxidase, and recovering the squalene epoxidase produced thereby.

ii. Sterol Methyl Transferase I

S-adenosyl-L-methionine:sterol C-24 methyl transferases (SMT1 and SMT2) catalyze the transfer of a methyl group from a cofactor, S-adenosyl-L-methionine, to the C-24 center of the sterol side chain (Bach, T.J. and Benveniste, P. (1997), Prog. Lipid Res. 36: 197-226). SMT in higher plant cells are responsible for their capability to produce a mixture of 24-methyl and 24-ethyl sterols (Schaffer, A., Bouvier-Navé, Benveniste, P., Schaller, H. (2000) Lipids 35: 263-269). Functional characterization of the SMT using a yeast *erg6* expression system demonstrated unambiguously that an SMT1 sequence encodes a cycloartenol-C24-methyltransferase and a SMT2 sequence encodes a 24-methylene lophenol-C24-methyltransferase in a given plant species (Bouvier-Navé, P., Husselstein, T., and Benveniste, P. (1998), Eur. J. Biochem. 246: 518-529). Several plant genes coding for SMT1 and SMT2 have been reported and reviewed (Schaffer, A., Bouvier-Navé, Benveniste, P., Schaller, H. (2000) Lipids 35: 263-

269). Transgenic plants expressing homologues of either SMT1 or SMT2 have been studied (Schaffer, A., Bouvier-Navé, Benveniste, P., Schaller, H. (2000) *Lipids* 35: 263-269). The use of these genes to modify plant sterol composition are also covered by two Monsanto patent applications (WO 98/45457 and WO 00/61771).

Sterol methyl transferase I enzymes known in the art are useful in the present invention. Exemplary sequences include the known *Arabidopsis* sterol methyl transferase I protein sequence Accession No. U71400 (disclosure SEQ ID NO:19), the known tobacco sterol methyl transferase I protein sequence Accession No. U81312 (disclosure SEQ ID NO:20) and *Ricinus communis* sterol-C-methyltransferase, *Eur. J. Biochem.*, **246**(2), 518-529 (1997). (Complete cds, Accession No. g2246457).

S-Adenosyl-L-Methionine-Sterol-C24-Methyltransferase--A nucleic acid sequence encoding an *Arabidopsis thaliana* S-adenosyl-L-methionine-sterol-C24-methyltransferase has been published by Husselstein et al. (1996) *FEBS Letters* 381: 87-92. Δ^{24} -sterol C-methyltransferase is enzyme number 2.1.1.41, Enzyme Nomenclature 1992, page 160.

25 iii. Sterol C4-Demethylase

Sterol C-4 demethylase catalyses the first of several demethylation reactions, which results in the removal of the two methyl groups at C-4. While in animals and fungi the removal of the two C-4 methyl groups occurs consecutively, in plants it has been reported that there are other steps between the first and second C-4 demethylations (Bach, T.J. and Benveniste, P. (1997), *Prog. Lipid Res.* 36: 197-226).

5

10

25

[illegible]

Sterol C-14 demethylase enzymes and sequences are known in the art. For example *Sorghum bicolor* obtusifolios 14 α -demethylase CYP51 mRNA, described in *Plant J.*, 11(2):191-201 (1997) (complete cds Accession No. U74319). In order to facilitate the modifications to sterol biosynthesis and accumulation described herein, the present invention also provides an isolated DNA molecule, having a nucleotide sequence selected from the group consisting of:

- 10 (a) obtusifolios C14 α -demethylase from clone ID: ATA101105 disclosure SEQ ID NO:14, clone ID ATA202967 disclosure SEQ ID NO:15, clone ID ATA403931 disclosure SEQ ID NO:17, or the complement thereof;
- (b) a nucleotide sequence that hybridizes to said
15 nucleotide sequence of (a) under a wash stringency equivalent to 0.5X SSC to 2X SSC, 0.1% SDS, at 55-65°C, and which encodes a polypeptide having obtusifolios C14 α -demethylase enzymatic activity substantially similar to that of the disclosed obtusifolios C14 α -
20 demethylase;
- (c) a nucleotide sequence encoding the same genetic information as said nucleotide sequence of (a), but which is degenerate in accordance with the degeneracy of the genetic code; and
- 25 (d) a nucleotide sequence encoding the same genetic information as said nucleotide sequence of (b), but which is degenerate in accordance with the degeneracy of the genetic code.

An additional aspect of the invention is the
30 recombinant constructs and vectors (pMON43842, Fig.29) comprising nucleic acid sequences encoding the novel obtusifolios C14 α -demethylase, as well as a method of producing the novel obtusifolios C14 α -demethylase,

T00290 62258850

[illegible]

Sterol C-5 desaturase catalyzes the insertion of the Δ^5 -double bond that normally occurs at the Δ^7 -sterol level, thereby forming a $\Delta^{5,7}$ -sterol (Parks et al., *Lipids* 30:227-230 (1995)). The reaction has been reported to involve the stereospecific removal of the 5 α and 6 α hydrogen atoms, biosynthetically derived from the 4 pro-R and 5 pro-S hydrogens of the (+) and (-) R-mevalonic acid, respectively (Goodwin, T.W. (1979) *Annu. Rev. Plant Physiol.* 30: 369-404). The reaction is obligatorily aerobic and requires NADPH or NADH. The desaturase has been reported to be a multienzyme complex present in microsomes. It consists of the desaturase itself, cytochrome b₅, and a pyridine nucleotide-dependent flavoprotein. The Δ^5 -desaturase is reported to be a mono-oxygenase that utilizes electrons derived from a reduced pyridine nucleotide via cytochrome b₅ (Taton, M., and Rahier, A. (1996) *Arch. Biochem. Biophys.* 325: 279-288). An *Arabidopsis thaliana* cDNA encoding a sterol-C5-desaturase was cloned by functional complementation of a yeast mutant, *erg3* defective in *ERG3*, the gene encoding the sterol C5-desaturase required for ergosterol biosynthesis (Gachotte D., Husselstein, T., Bard, M., Lacroute F., and Benveniste, P. (1996) *The Plant Journal* 9(3): 391-398). Known sterol C5-desaturase enzymes are useful in the present invention, including *Arabidopsis* sterol C5-

desaturase protein sequence Accession No. X90454,
disclosure SEQ ID NO:22, and the *Arabidopsis thaliana*
mRNA for sterol-C5-desaturase described in *Plant J.*
9(3):391-398 (1996) (complete cds Accession No.
5 g1061037).

The NCBI (National Center for Biotechnology
Information) database shows 37 sequences for sterol
desaturase that are useful in the present invention.
The following are exemplary of such sequences. From
10 yeast: C5 sterol desaturase NP_013157 (*Saccharomyces*
cerevisiae); hypothetical C5 sterol desaturase-fission
T40027 (*Schizosaccharomyces pombe*); C5 sterol
desaturase-fission T37759 (*Schizosaccharomyces pombe*);
C5 sterol desaturase JQ1146 (*Saccharomyces cerevisiae*);
15 C5 sterol desaturase BAA21457 (*schizosaccharomyces*
pombe); C5 sterol desaturase CAA22610
(*Schizosaccharomyces pombe*); putative C5 sterol
desaturase CAA16898 (*Schizosaccharomyces pombe*);
probable C5 sterol desaturase O13666 (*erg3 _schpo*); C5
20 sterol desaturase P50860 (*Erg3_canga*); C5 sterol
desaturase P32353 (*erg3_yeast*); C5,6 desaturase
AAC99343 (*Candida albicans*); C5 sterol desaturase
BAA20292 (*Saccharomyces cerevisiae*); C5 sterol
desaturase AAB39844 (*Saccharomyces cerevisiae*); C5
25 sterol desaturase AAB29844 (*Saccharomyces cerevisiae*);
C5 sterol desaturase CAA64303 (*Saccharomyces*
cerevisiae); C5 sterol desaturase AAA34595
(*Saccharomyces cerevisiae*); C5 sterol desaturase
AAA34594 (*Saccharomyces cerevisiae*). From plants: C5
30 sterol desaturase S71251 (*Arabidopsis thaliana*);
putative sterol-C5-desaturase AAF32466 (*Arabidopsis*
thaliana); sterol-C5-desaturase AAF32465 (*Arabidopsis*
thaliana); putative sterol desaturase AAF22921
(*Arabidopsis thaliana*); delta7 sterol C5 desaturase

T00290 225860

[illegible]

25 vi. Sterol Methyl Transferase II

Known sterol methyl transferase II enzymes are useful in the present invention, including *Arabidopsis* sterol methyl transferase II protein sequence (complete

Recombinant constructs encoding any of the foregoing enzymes affecting the steroid biosynthetic pathway can be incorporated into recombinant vectors comprising the recombinant constructs comprising the isolated DNA molecules. Such vectors can be bacterial or plant expression vectors.

The present invention contemplates a recombinant construct that contains a DNA sequence encoding a polypeptide exhibiting 3-hydroxy-3-methylglutaryl-Coenzyme A (HMG-CoA) reductase activity and a DNA sequence encoding a polypeptide exhibiting the activity of a steroid pathway enzyme. Each polypeptide-encoding DNA sequence is operably linked in the 5' to 3' direction independent of the other sequence. Each DNA sequence in the 5' to 3' direction comprises a promoter, then the DNA sequence encoding the polypeptide then a transcription termination signal sequence. The steroid pathway enzyme is a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifolioside C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, or a sterol methyl transferase II enzyme.

Preferably, the promoters in the recombinant construct are seed-specific promoters. In one embodiment, the promoter is derived from a species in a different order from the host cell. In other 30 embodiments, the encoded HMG CoA reductase and/or steroid pathway enzymes is(are) from a species in a different order from the order that of the host cell.

[illegible]

Useful in expressing the polypeptide coding gene are promoters that are inducible, viral, synthetic, constitutive as described by Poszkowski et al., *EMBO J.*, **3**:2719 (1989) and Odell et al., *Nature*, **313**:810 (1985), and temporally regulated, spatially regulated and spatiotemporally regulated as given in Chau et al., *Science*, **244**:174-181 (1989). The promoter preferably comprises a promoter sequence whose function in regulating expression of the structural gene is substantially unaffected by the amount of sterol or squalene in the cell. As used herein, the term "substantially unaffected" means that the promoter is

5

15

25

30

A plant integrating vector containing a structural gene coding for a polypeptide having HMG-CoA reductase activity is engineered to be under control of the Lectin promoter and that vector is introduced into soybean plants using a protoplast transformation method. E.G. Dhir et al., *Plant Cell Reports*, 10:97 (1991). The expression of the polypeptide having HMG-CoA reductase activity is directed specifically to the seeds of the transgenic plant. In this way, a transgenic soybean seed having increased squalene accumulation is produced. Such seeds can then be used to extract oil containing enhanced levels of squalene. As set forth hereinafter, such squalene-enhanced oil is characterized by a greater thermal stability when compared to non-squalene-enhanced oil.

In the present invention, a plant has an exogenously provided structural gene for HMG-CoA reductase and at least one of the six enumerated steroid pathway enzymes, a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifoliol C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, or a sterol methyl transferase II enzyme. The plant or seed thus containing these added genes is contemplated, while the methods to arrive at a plant or seed according to the invention are open to the multitude of methods contemplated by a person of ordinary skill in the art. In particular, all of the added structural genes do not have to have been added at the same time, or by the same route. Thus, for example, the HMG-CoA reductase activity may result from a cross with a plant made according to a process of U.S. Patent No. 5,349,126, while a steroid pathway enzyme is added by nucleic acid

bombardment to that plant. Further, when more than one nucleotide sequence encoding a steroid pathway enzyme is present in a contemplated plant, the expression of the gene does not have to be under the control of the same promoter, or even the same type of promoter.

The choice of which plant integrating vector and ultimately to which promoter a polypeptide coding gene is operatively linked depends directly on the functional properties desired, e.g. the location and timing of protein expression, and the host cell to be transformed. These are well known limitations inherent in the art of constructing recombinant DNA molecules. However, a vector useful in practicing the present invention is capable of directing the expression of the polypeptide coding gene, i.e., the gene encoding HMG-CoA reductase activity, included in the DNA segment to which it is operatively linked.

Typical vectors useful for expression of genes in higher plants are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens*, described by Rogers et al., *Meth. Enzymol.*, **153**:253-277 (1987). However, several other plant integrating vector systems are known to function in plants including pCaMVCN transfer control vector described by Fromm et al. *Proc. Nat. Acad. Sci. USA*, **82**:5824 (1985). Plasmid pCaMVCN (available from Pharmacia, Piscataway, N.J.) includes the cauliflower mosaic virus CaMV 35S promoter.

The use of retroviral plant integrating vectors to form the recombinant DNAs of the present invention is also contemplated. As used herein, the term "retroviral plant integrating vector" refers to a DNA molecule that includes a promoter sequence derived from

the long terminal repeat (LTR) region of a retrovirus genome.

In preferred embodiments, the vector used to express the polypeptide coding gene includes a selection marker that is effective in a plant cell, preferably a drug resistance selection marker. One preferred drug resistance marker is the gene whose expression results in kanamycin resistance; i.e., the chimeric gene containing the nopaline synthase promoter, Tn5 neomycin phosphotransferase II and nopaline synthase 3' nontranslated region described by Rogers et al., in *Methods for Plant Molecular Biology*, A. Weissbach and H. Weissbach eds., Academic Press Inc., San Diego, Calif. (1988).

A variety of methods have been developed to operatively link DNA to vectors via complementary cohesive termini or blunt ends. For instance, complementary homopolymer tracts can be added to the DNA segment to be inserted and to the vector DNA. The vector and DNA segment are then joined by hydrogen bonding between the complementary homopolymeric tails to form recombinant DNA molecules.

Alternatively, synthetic linkers containing one or more restriction endonuclease sites can be used to join the DNA segment to the plant integrating vector. The synthetic linkers are attached to blunt-ended DNA segments by incubating the blunt-ended DNA segments with a large excess of synthetic linker molecules in the presence of an enzyme that is able to catalyze the ligation of blunt-ended DNA molecules, such as bacteriophage T4 DNA ligase. Thus, the products of the reaction are DNA segments carrying synthetic linker sequences at their ends. These DNA segments are then cleaved with the appropriate restriction endonuclease

100290-6253660

5 endonuclease sites are commercially available from a
number of sources including New England BioLabs,
Beverly, Mass.

Also contemplated by the present invention are RNA equivalents of the above-described recombinant DNA molecules.

A preferred recombinant DNA molecules utilized in accordance with the present invention are pMON53733-pMON53740 (Figures 13-20).

15 A. Promoters and Target Tissues

Promoters useful in the present invention include those that confer appropriate cellular and temporal specificity of expression. Such promoters include those that are constitutive or inducible, environmentally- or developmentally-regulated, or organelle-, cell-, or tissue-specific. Preferred promoters for use with the present invention promote expression of the introduced enzymes in the seed in the cytosol, although expression in plasmids or organelles of the seeds is also contemplated.

Often-used constitutive promoters include the CaMV 35S promoter (Odell et al. (1985) *Nature* 313: 810), the enhanced CaMV 35S promoter, the Figwort Mosaic Virus (FMV) promoter (Richins et al. (1987) *NAR* 20: 8451),
30 the mannopine synthase (*mas*) promoter, the nopaline synthase (*nos*) promoter, and the octopine synthase (*ocs*) promoter.

Useful inducible promoters include heat-shock promoters (Ou-Lee et al. (1986) *Proc. Natl. Acad. Sci.*

USA 83: 6815; Ainley et al. (1990) *Plant Mol. Biol.* 14: 949), a nitrate-inducible promoter derived from the spinach nitrite reductase gene (Back et al. (1991) *Plant Mol. Biol.* 17: 9), hormone-inducible promoters
5 (Yamaguchi-Shinozaki et al. (1990) *Plant Mol. Biol.* 15: 905; Kares et al. (1990) *Plant Mol. Biol.* 15: 905), and light-inducible promoters associated with the small subunit of RuBP carboxylase and LHCP gene families (Kuhlemeier et al. (1989) *Plant Cell* 1: 471; Feinbaum
10 et al. (1991) *Mol. Gen. Genet.* 226: 449; Weisshaar et al. (1991) *EMBO J.* 10: 1777; Lam and Chua (1990) *Science* 248: 471; Castresana et al. (1988) *EMBO J.* 7: 1929; Schulze-Lefert et al. (1989) *EMBO J.* 8: 651).

Examples of useful tissue-specific,
15 developmentally-regulated promoters include fruit-specific promoters such as the E4 promoter (Cordes et al. (1989) *Plant Cell* 1:1025), the E8 promoter (Deikman et al. (1988) *EMBO J.* 7: 3315), the kiwifruit actinidin promoter (Lin et al. (1993) *PNAS* 90: 5939), the 2A11
20 promoter (Houck et al., U.S. Patent 4,943,674), and the tomato pZ130 promoter (U.S. Patents 5,175, 095 and 5,530,185); the β -conglycinin 7S promoter (Doyle et al. (1986) *J. Biol. Chem.* 261: 9228; Slighton and Beachy (1987) *Planta* 172: 356), and seed-specific promoters
25 (Knutzon et al. (1992) *Proc. Natl. Acad. Sci. USA* 89: 2624; Bustos et al. (1991) *EMBO J.* 10: 1469; Lam and Chua (1991) *J. Biol. Chem.* 266: 17131; Stayton et al. (1991) *Aust. J. Plant. Physiol.* 18: 507). Fruit-specific gene regulation is discussed in U.S. Patent
30 5,753,475. Other useful seed-specific promoters include, but are not limited to, the napin, phaseolin, zein, soybean trypsin inhibitor, 7S, ADR12, ACP, stearyl-ACP desaturase, oleosin, *Lasquerella*

5 include the corn globulin 1 and oleosin promoters. Useful endosperm-specific promoters include the rice glutelin-1 promoter, the promoters for the low-pI -amylase gene (Amy32b) (Rogers et al. (1984) *J. Biol. Chem.* 259: 12234), the high-pI -amylase gene (Amy 64) (Khurseed et al. (1988) *J. Biol. Chem.* 263: 18953), and
10 the promoter for a barley thiol protease gene ("Aleurain") (Whittier et al. (1987) *Nucleic Acids Res.* 15: 2515).

15 production of sterol compounds such as sitosterol,
sitosterol esters, sitostanol, sitostanol esters, and
tocopherols, and reduced production of campesterol,
campestanol, and esters thereof, include, but are not
20 limited to, fruits, flowers, seeds, roots, tubers,
leaves, stems, buds, and other vegetable parts of
plants. Within seeds, appropriate organ compartments
include the embryo, the endosperm, and the aleurone
layer. Within any of the noted target tissues,
appropriate cellular compartments include, but are not
25 limited to, the cell cytoplasm and plastids (e.g.,
proplastids, chloroplasts, chromoplasts, leucoplasts,
amyloplast, etc.).

B. Vectors

30 In plants, transformation vectors capable of introducing encoding DNAs involved in sterol compound and tocopherol biosynthesis are easily designed, and generally contain one or more DNA coding sequences of interest under the transcriptional control of 5' and 3'

regulatory sequences. Such vectors generally comprise, operatively linked in sequence in the 5' to 3' direction, a promoter sequence that directs the transcription of a downstream heterologous structural DNA in a plant; optionally, a 5' non-translated leader sequence; a nucleotide sequence that encodes a protein of interest; and a 3' non-translated region that encodes a polyadenylation signal which functions in plant cells to cause the termination of transcription and the addition of polyadenylate nucleotides to the 3' end of the mRNA encoding the protein. Plant transformation vectors also generally contain a selectable marker. Typical 5'-3' regulatory sequences include a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal. Vectors for plant transformation have been reviewed in Rodriguez et al. (1988) *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, Butterworths, Boston; Glick et al. (1993) *Methods in Plant Molecular Biology and Biotechnology* CRC Press, Boca Raton, Fla; and Croy (1993) In *Plant Molecular Biology Labfax*, Hames and Rickwood, Eds., BIOS Scientific Publishers Limited, Oxford, UK.

The use of transit peptides, e.g. translational fusion peptides, are not preferred for use in conjunction with the enzymes of the present invention, where the sterol synthethic compounds are present primarily in the cellular cytosol.

V. Cell Transformation and Plant Regeneration

The amount of a gene coding for a polypeptide having HMG-CoA reductase activity is increased by

transforming a desired plant cell with a suitable vector that contains that added exogenous structural gene. Expression of that gene in the transformed plant cell and transgenic plants developed from that transformed plant cell enhances the activity of HMG-CoA reductase.

Methods for transforming polypeptide-coding genes into plant cells include *Agrobacterium*-mediated plant transformation, protoplast transformation, gene transfer into pollen, injection into reproductive organs and injection into immature embryos. Each of these methods has distinct advantages and disadvantages. Thus, one particular methods of introducing genes into a particular plant strain may not necessarily be the most effective for another plant strain, but it is well known which methods are useful for a particular plant strain.

Agrobacterium-mediated transfer is a widely applicable system for introducing genes into plant cells because the DNA can be introduced into whole plant tissues, thereby bypassing the need for regeneration of an intact plant from a protoplast. The use of *Agrobacterium*-mediated plant integrating vectors to introduce DNA into plant cells is well known in the art. See, for example, the methods described by Fraley et al., *Biotechnology*, 3:629 (1984) and Rogers et al., *Methods in Enzymology*, 153:253-277 (1987). Further the integration of the T8-DNA is a relatively precise process resulting in few rearrangements. The region of DNA to be transferred is defined by the border sequences, and intervening DNA is usually inserted into the plant genome as described by Spielmann et al., *Mol. Gen. Genet.*, 205:34 (1986) and Jorgensen et al, *Mol. Gen. Genet.*, 207:471 (1987).

09885723.062001

Modern *Agrobacterium* transformation vectors are capable of replication in *E. coli* as well as *Agrobacterium*, allowing for convenient manipulations as described by Klee et al., in *Plant DNA Infectious Agents*, T. Hohn and J. Schell, eds., Springer-Verlag, New York (1985) pp. 179-203.

Moreover, recent technological advances in vectors for *Agrobacterium*-mediated gene transfer have improved the arrangement of genes and restriction sites in the vectors to facilitate construction of vectors capable of expressing various polypeptide coding genes. The vectors described by Rogers et al., *Meth. Enzymol.*, 153:253 (1987), have convenient multi-linker regions flanked by a promoter and a polyadenylation site for directed expression inserted polypeptide coding genes and are suitable for present purposes.

In addition, *Agrobacteria* containing both armed and disarmed Ti genes can be used for the transformations. Both types of transforming systems are illustrated herein. Transformants from the former system result in callus from which the desired squalene or sterol can be obtained, whereas transformants obtained from the latter, disarmed Ti genes can be regenerated into complete transgenic plants from whose tissues, e.g. leaf, seed and root, the desired chemicals can be obtained.

In those plant strains where *Agrobacterium*-mediated transformation is efficient, it is the method of choice because of the facile and defined nature of the gene transfer.

Agrobacterium-mediated transformation of leaf disks and other tissues, such as cotyledons and hypocotyls, appears to be limited to plant strains that *Agrobacterium* naturally infects. *Agrobacterium*-

09885723-062004
T00290" E2Z58860

5

15

30

segregant transgenic plant that contains the added genes according to the invention, germinating some of the seed produced and analyzing the resulting plants produced for enhanced HMG-CoA reductase activity, 5 steroid pathway product accumulation or both, relative to a control (native, non-transgenic) or an independent segregant transgenic plant.

A homozygous transgenic plant exhibits enhanced HMG-CoA reductase activity as compared to a native, 10 non-transgenic plant and an independent segregant transgenic plant.

It is to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating 15 added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes that encode a polypeptide having HMG-CoA reductase activity. Back-crossing to a parental plant and outcrossing with a non-transgenic plant are 20 also contemplated.

A. Host Cells and Transformed Plant Cells

Cells modified according to the present invention are contemplated at each stage of the processes of the 25 invention. As a result of the invention comprising at least two genes, there are several means to accomplish that end. In some embodiments of the invention, the intermediate steps include transformation of nucleic acids comprising some or all of the genes into host 30 cells.

The nucleic acid sequence encoding a polypeptide exhibiting HMGR activity does not have to be in the same orientation as a nucleic acid sequence encoding a polypeptide exhibiting the activity of a steroid

pathway enzyme. The coding nucleic acids can be under the control of different promoters or be in different orientations. For the host plant cell useful in carrying out the steroid compound biosynthesis

5 according to the invention, the minimum that is required is the coding nucleic acids be in the same host cell. As long as the HMGR and a steroid pathway enzyme coding sequences are present in the same host cell, they do not have to be on the same DNA molecule

10 or under the control of the same promoter, nor do they have to be derived from the same vector or construct.

Host cells are useful for making, storing, reproducing or manipulating nucleic acid constructs of the invention. Contemplated host cells are eukaryotic

15 cells, such as yeast or plant cells. Any plant cells can be utilized with the present invention. Some particularly useful agriculturally significant plant cells are canola, soybean, corn, maize, tobacco, cotton, rape, tomato and alfalfa. Other common plant

20 varieties are carrot, barley, arabidopsis, guayule and petunia. Prokaryotic host cells containing constructs and/or vectors according to the invention are also contemplated (e.g. *E. coli*).

One embodiment of the invention is a transformed

25 host cell containing inter alia a recombinant construct that encodes both a DNA sequence encoding a polypeptide exhibiting 3-hydroxy-3-methylglutaryl-Coenzyme A reductase enzyme activity and a DNA sequence encoding a steroid pathway enzyme, where the steroid pathway

30 enzyme is as described in detail above. In a preferred embodiment, those coding DNA sequences are operably linked to a promoter and a transcription termination signal sequence. In the coding sense direction of the construct, the components of the construct are operably

098872-02001
"00290" 02/28/90

Another embodiment of the invention is host cell
5 that has been transformed with a recombinant vector
that has such a construct. As discussed herein, in one
embodiment of the invention, such a recombinant vector
is a plant expression vector. Preferably such a host
cell is a plant cell.

Non-limiting examples of plants that can be used in the practice of the invention include, acacia, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beans, beet, blackberry, blueberry, broccoli, brussel sprouts, cabbage, canola, cantaloupe, carrot, cassava, cauliflower, celery, cherry, chicory, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, figs, garlic, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, mushroom, nectarine, nut, oat, oil palm, oil seed rape, okra, onion, orange, ornamental plants, papaya, parsley, pea, peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet

potato, sweetgum, tangerine, tea, tobacco, tomato, triticale, turf, turnip, watermelon, wheat, yam, and zucchini.

Plants particularly attractive for the steroid pathway modifications described herein include those that produce carbon substrates which can be employed for synthesis of these compounds. Non-limiting examples of such plants include various monocots and dicots, including high oil seed plants such as high oil seed *Brassica* (e.g., *Brassica nigra*, *Brassica napus*, *Brassica hirta*, *Brassica rapa*, *Brassica campestris*, *Brassica carinata*, and *Brassica juncea*), soybean (*Glycine max*), castor bean (*Ricinus communis*), cotton, safflower (*Carthamus tinctorius*), sunflower (*Helianthus annuus*), flax (*Linum usitatissimum*), corn (*Zea mays*), coconut (*Cocos nucifera*), palm (*Elaeis guineensis*), oilnut trees such as olive (*Olea europaea*), sesame, and peanut (*Arachis hypogaea*), as well as *Arabidopsis*, tobacco, wheat, barley, oats, amaranth, potato, rice, tomato, and legumes (e.g., peas, beans, lentils, alfalfa, etc.).

Enhancement of sitostanol compound production by the methods discussed herein is expected to result in yields of sitostanol, sitostanol esters, or mixtures thereof in an amount of at least about 57% by weight, preferably from about 57% to about 90% by weight, and more preferably from about 57% to about 65% by weight of the total sterol compounds present in seed oil. Expressed on a seed dry weight basis, sitostanol, sitostanol esters, or mixtures thereof are expected to be present in an amount of at least about 0.08%, preferably from about 0.08% to about 0.8%, and more preferably from about 0.08% to about 0.4% of seed dry weight.

[illegible]

meristems, etc., to generate transgenic plants. These methods include, for example, *Agrobacterium*-mediated transformation, particle gun delivery, microinjection, electroporation, polyethylene glycol-mediated protoplast transformation, liposome-mediated transformation, etc. (reviewed in Potrykus (1991) *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42: 205).

15 compound and tocopherol biosynthesis and accumulation described herein can be produced by transforming plant cells with a DNA construct as described above via any of the foregoing methods; selecting plant cells that have been transformed on a selective medium;

20 regenerating plant cells that have been transformed to produce differentiated plants; and selecting a transformed plant that expresses the enzyme-encoding nucleotide sequence(s) at a level such that the amount of sitosterol, sitosterol esters, sitostanol,

25 sitostanol esters, tocopherol compound(s), and campesterol/campestanol and their esters is within the ranges described herein.

on the same vector), a co-transformation event (all necessary DNAs present on separate vectors that are introduced into plants or plant cells simultaneously), or by independent transformation events (all necessary DNAs present on separate vectors that are introduced

into plants or plant cells independently). Traditional breeding methods can subsequently be used to incorporate the desired combination of enzymes into a single plant, and to produce hybrid progeny of the invention plants.

Specific methods for transforming a wide variety of dicots and obtaining transgenic plants are well documented in the literature (Gasser and Fraley (1989) *Science* 244: 1293; Fisk and Dandekar (1993) *Scientia Horticulturae* 55: 5; Christou (1994) *Agro Food Industry Hi Tech*, p. 17; and the references cited therein).

Apomixis is a genetically controlled method of reproduction in plants where the embryo is formed without union of an egg and a sperm. There are three basic types of apomictic reproduction: 1) apospory where the embryo develops from a chromosomally unreduced egg in an embryo sac derived from the nucellus, 2) diplospory where the embryo develops from an unreduced egg in an embryo sac derived from the megaspore mother cell, and 3) adventitious embryony where the embryo develops directly from a somatic cell. In most forms of apomixis, pseudogamy or fertilization of the polar nuclei to produce endosperm is necessary for seed viability. In apospory, a "nurse" cultivar can be used as a pollen source for endosperm formation in seeds. The nurse cultivar does not affect the genetics of the aposporous apomictic cultivar since the unreduced egg of the cultivar develops parthenogenetically, but makes possible endosperm production.

Apomixis is economically important, especially in transgenic plants, because it causes any genotype, no matter how heterozygous, to breed true. Thus, with apomictic reproduction, heterozygous transgenic plants

can maintain their genetic fidelity throughout repeated life cycles. Methods for the production of apomictic plants are known in the art. See, U.S. Patent No. 5,811,636 and references cited therein which are herein
5 incorporated by reference.

The present invention also encompasses uniform populations of any of the plants discussed herein.

Successful transformation and plant regeneration have been achieved in the monocots as follows:

10 asparagus (*Asparagus officinalis*; Bytebier et al. (1987) *Proc. Natl. Acad. Sci. USA* 84: 5345); barley (*Hordeum vulgare*; Wan and Lemaux (1994) *Plant Physiol.* 104: 37); maize (*Zea mays*; Rhodes et al. (1988) *Science* 240: 204; Gordon-Kamm et al. (1990) *Plant Cell* 2: 603;
15 Fromm et al. (1990) *Bio/Technology* 8: 833; Koziel et al. (1993) *Bio/Technology* 11: 194); oats (*Avena sativa*; Somers et al. (1992) *Bio/Technology* 10: 1589); orchardgrass (*Dactylis glomerata*; Horn et al. (1988) *Plant Cell Rep.* 7: 469); rice (*Oryza sativa*, including
20 indica and japonica varieties; Toriyama et al. (1988) *Bio/Technology* 6: 10; Zhang et al. (1988) *Plant Cell Rep.* 7: 379; Luo and Wu (1988) *Plant Mol. Biol. Rep.* 6: 165; Zhang and Wu (1988) *Theor. Appl. Genet.* 76: 835; Christou et al. (1991) *Bio/Technology* 9: 957); rye
25 (*Secale cereale*; De la Pena et al. (1987) *Nature* 325: 274); sorghum (*Sorghum bicolor*; Cassas et al. (1993) *Proc. Natl. Acad. Sci. USA* 90: 11212); sugar cane (*Saccharum* spp.; Bower and Birch (1992) *Plant J.* 2: 409); tall fescue (*Festuca arundinacea*; Wang et al.
30 (1992) *Bio/Technology* 10: 691); turfgrass (*Agrostis palustris*; Zhong et al. (1993) *Plant Cell Rep.* 13: 1); and wheat (*Triticum aestivum*; Vasil et al. (1992) *Bio/Technology* 10: 667; Weeks et al. (1993) *Plant*

09085723.062004
T00290" 22/58860

Physiol. 102: 1077; Becker et al. (1994) *Plant J.* 5: 299).

Plant transformation vectors capable of delivering DNAs (genomic DNAs, plasmid DNAs, cDNAs, or synthetic DNAs) encoding plant-derived or other enzymes that affect the biosynthesis and accumulation of sterol compounds and tocopherols in plants for optimizing the pools of sitosterol, sitostanol, esters of either, and tocopherols, and for reducing the levels of campesterol, campestanol, and/or their esters, can be easily designed by art-recognized methods. Various strategies can be employed to introduce these encoding DNAs into plants to produce transgenic plants that biosynthesize and accumulate desirable levels of various sterol compounds and tocopherols, including:

1. Transforming individual plants with an encoding DNA of interest. Two or more transgenic plants, each containing one of these DNAs, can then be grown and cross-pollinated so as to produce hybrid plants containing the two DNAs. The hybrid can then be crossed with the remaining transgenic plants in order to obtain a hybrid plant containing all DNAs of interest within its genome.

2. Sequentially transforming plants with plasmids
25 containing each of the encoding DNAs of interest,
respectively.

3. Simultaneously cotransforming plants with plasmids containing each of the encoding DNAs, respectively.

30 4. Transforming plants with a single plasmid
containing two or more encoding DNAs of interest.

5. Transforming plants by a combination of any of the foregoing techniques in order to obtain a plant

that expresses a desired combination of encoding DNAs of interest.

Traditional breeding of transformed plants produced according to any one of the foregoing methods by successive rounds of crossing can then be carried out to incorporate all the desired encoding DNAs in a single homozygous plant line (Nawrath et al. (1994) *Proc. Natl. Acad. Sci. USA* 91: 12760; PCT International Publication WO 93/02187), or to produce hybrid offspring.

In methods 2 and 3, the use of vectors containing different selectable marker genes to facilitate selection of plants containing two or more different encoding DNAs is advantageous. Examples of useful selectable marker genes include those conferring resistance to kanamycin, hygromycin, sulphonamides, glyphosate, bialaphos, and phosphinothricin.

C. Processes of Regeneration

Processes of regeneration of plants from transformed protoplasts are known in the art.

D. Transgenic Plants and Progeny

The present invention contemplates the plants that contain the exogenous constructs according to the present invention, such that a plant comprises at least one transformed plant cell comprising a nucleic acid construct. The nucleic acid construct, as described above has as operably linked components in the 5' to 3' direction, a promoter, a DNA sequence encoding a polypeptide exhibiting 3-hydroxy-3-methylglutaryl-Coenzyme A reductase enzyme activity, and a transcription termination signal sequence. The plant also comprises a nucleic acid construct that has as

Variable	Mean	SD	Min	Max
Age	34.5	10.2	21	55
Gender	Male	Female		
Marital status	Married	Single		
Education	High school	College		
Occupation	Manager	Worker		
Income	Low	High		
Health status	Good	Poor		
Stress level	Low	High		
Life satisfaction	Low	High		
Resilience	Low	High		
Optimism	Low	High		
Self-efficacy	Low	High		
Perceived social support	Low	High		
Perceived stress	Low	High		
Depression	Low	High		
Anxiety	Low	High		
Quality of life	Low	High		
Health-related quality of life	Low	High		
Physical health	Low	High		
Mental health	Low	High		
Overall health	Low	High		
Life expectancy	Low	High		
Healthcare utilization	Low	High		
Health insurance	Low	High		
Healthcare access	Low	High		
Healthcare quality	Low	High		
Healthcare cost	Low	High		
Healthcare satisfaction	Low	High		
Healthcare utilization	Low	High		
Healthcare access	Low	High		
Healthcare quality	Low	High		
Healthcare cost	Low	High		
Healthcare satisfaction	Low	High		

The present invention contemplates a plant, the genome of which comprises introduced DNA. That introduced DNA has at least two components. One component is a DNA encoding a 3-hydroxy-3-methylglutaryl-Coenzyme A reductase enzyme. The other component is DNA encoding a steroid pathway enzyme that is a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifoliol C14 α -demethylase enzyme, a sterol C5-desaturase enzyme, or a sterol methyl transferase II

enzyme. The storage organs, preferably seeds, of such a plant contain an elevated level of total accumulated sterol, compared to storage organs of an otherwise identical plant, the genome of which does not comprise the introduced DNA. The introduced DNAs are
5 operatively linked to regulatory signals, preferably that cause seed-specific expression of said introduced DNAs. The seeds of such a plant contain a reduced level of squalene, cycloartenol, 24-methylene
10 cycloartenol, obtusifoliol, stigmasta-7-enol, or campesterol compared to the seeds of an otherwise identical plant or compared to a plant comprising an introduced DNA encoding a HMG CoA reductase enzyme without the contemplated steroid pathway enzyme.

15 Also contemplated is a plant with introduced DNA, as described above, that produces seed having an elevated level of a steroid pathway product, compared to a corresponding transgenic or non-transgenic plant that does not contain said introduced DNA.

20 The invention also contemplates a plant comprising introduced DNA encoding (i) an HMGR enzyme and (ii) a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifoliol C14 α -demethylase enzyme, a sterol C5-
25 desaturase enzyme, a sterol methyl transferase II enzyme, or mixtures thereof, wherein said plant that produces a storage organ (preferably a seed) having an elevated level of a sterol pathway product compared to a corresponding transgenic or non-transgenic plant that
30 does not contain said introduced DNA.

The invention also contemplates a plant having introduced DNA, as described above, that produces a storage organ (preferably a seed) having a reduced

09865729.062004
T00290"E2Z5B850

THE **NEW** **YORK** **PUBLIC** **LIBRARY**

15

20

25

30

Another embodiment is a seed resulting from a cross of a plant having introduced DNA, described above, with a nurse cultivar. Also contemplated are seeds of any of the above-described plants. Also part
5 of the present invention plant parts, other than a seed of any of the above-described plants.

Uniform populations of the above-described plants are also contemplated.

10 E. Stability of Transgene Expression

As several overexpressed enzymes may be required to produce optimal levels of substrates for the biosynthesis of sterol compounds and tocopherols, the phenomenon of co-suppression may influence transgene
15 expression in transformed plants. Several strategies can be employed to avoid this potential problem (Finnegan and McElroy (1994) *Bio/Technology* 12: 883).

One commonly employed approach is to select and/or screen for transgenic plants that contain a single
20 intact copy of the transgene or other encoding DNA (Assaad et al. (1993) *Plant Mol. Biol.* 22: 1067; Vaucheret (1993) *C.R. Acad. Sci. Paris, Science de la vie/Life Sciences* 316: 1471; McElroy and Brettell (1994) *TIBTECH* 12: 62). *Agrobacterium*-mediated
25 transformation technologies are preferred in this regard.

Inclusion of nuclear scaffold or matrix attachment regions (MAR) flanking a transgene has been shown to increase the level and reduce the variability
30 associated with transgene expression in plants (Stief et al. (1989) *Nature* 341: 343; Breyne et al. (1992) *Plant Cell* 4: 463; Allen et al. (1993) *Plant Cell* 5: 603); Mlynarova et al. (1994) *Plant Cell* 6: 417; Spiker and Thompson (1996) *Plant Physiol.* 110: 15). Flanking a

098857-062004
"E2258350"

transgene or other encoding DNA with MAR elements may overcome problems associated with differential base composition between such transgenes or encoding DNAs and integrations sites, and/or the detrimental effects of sequences adjacent to transgene integration sites.

The use of enhancers from tissue-specific or developmentally-regulated genes may ensure that expression of a linked transgene or other encoding DNA occurs in the appropriately regulated manner.

The use of different combinations of promoters, plastid targeting sequences, and selectable markers for introduced transgenes or other encoding DNAs can avoid potential problems due to *trans*-inactivation in cases where pyramiding of different transgenes within a single plant is desired.

Finally, inactivation by co-suppression can be avoided by screening a number of independent transgenic plants to identify those that consistently overexpress particular introduced encoding DNAs (Register et al. (1994) *Plant Mol. Biol.* 25: 951). Site-specific recombination in which the endogenous copy of a gene is replaced by the same gene, but with altered expression characteristics, should obviate this problem (Yoder and Goldsbrough (1994) *Bio/Technology* 12: 263).

Any of the foregoing methods, alone or in combination, can be employed in order to insure the stability of transgene expression in transgenic plants of the present invention.

F. Hybrid Plants

The invention contemplates a plant having introduced DNA encoding an HMGR and at least one of the six steroid pathway enzymes, as described in detail above. It is contemplated that a transgenic plant

having DNA encoding an HMGR, as is known in the art, might be crossed with a transgenic plant having DNA encoding at least one of the six steroid pathway enzymes.

5 Also contemplated as a hybrid plant according to the invention is a plant that is a hybrid of a transgenic plant having introduced DNA encoding an HMGR and at least one of the six steroid pathway enzymes wherein the plant has been hybridized with another
10 strain, yet still retains the introduced DNA.

G. Storage Organs

The term "storage organ" as used herein, refers to the seeds, fruits or vegetable parts of a plant. Most
15 often the seed is important for use in the present invention. However, there are consumable embodiments, such as with potatoes or carrots, where the vegetable parts are preferred.

A contemplated embodiment of the present invention
20 is a storage organ comprising at least one transformed host cell. The transformed host cell has at a minimum a construct according the invention as described above. Also contemplated are the embodiments when the construct has plant promoters, when the construct is
25 recombinant, when the construct is part of a vector, and when the vector is a plant expression vector.

The invention contemplates a transgenic plant seed transformed with a vector comprising a DNA segment that encodes a polypeptide having HMG-CoA reductase
30 activity, and a DNA segment that encodes a polypeptide having a steroid pathway enzyme, wherein the transgenic plant seed is capable of germinating into a transgenic plant that over-accumulates steroid pathway products relative to a native, non-transgenic plant of the same

09885723.062004
"00290" 22/58850

[illegible]

10

20

30

[illegible]

10

15

VI. Harvest

25

30

[illegible]

Oils produced by the cells, plants, and methods disclosed herein are superior in
25 phytosterol/phytostanol composition to conventional oils in a variety of ways. Oil of the present invention can contain an elevated level of at least one sterol, at least one phytosterol, at least one phytosterol ester, at least one phytostanol, at least
30 one phytostanol ester, or mixtures thereof. Preferred compounds include sitosterol, sitostanol, and their esters.

Oil from seed of plants containing and expressing introduced DNA encoding a sterol methyltransferase II

enzyme advantageously contains a reduced level of campesterol, at least one campesterol ester, campestanol, at least one campestanol ester, or mixtures thereof. The sterol methyltransferase II-
5 encoding DNA can be introduced alone, or in combination with other introduced DNA sequences encoding enzymes affecting the biosynthesis of steroid compounds as discussed herein. Campesterol/campestanol and their esters are considered to be undesirable because they
10 are readily absorbed in the intestine, while their safety in the blood is unknown. Employing the plants and methods disclosed herein, one can obtain seed oil comprising about 0% to about 19%, preferably about 0% to about 12%, more preferably about 5% to about 9%
15 campesterol, at least one campesterol ester, campestanol, at least one campestanol ester, or mixtures thereof by weight of the total sterol compounds of the oil. (The levels of these compounds are difficult to express on a percent seed dry weight
20 basis because different seeds contain different percentages of these compounds expressed on this basis) These values represent a reduction of about 10% to about 100% in the amount of these compounds compared to those in conventional oils.

25 Introduction into plant cells of the enzyme-encoding DNA sequences discussed above modifies the biosynthesis of sterol compounds carried out by the methods, and in the cells, plants, and seeds, disclosed herein. In particular, the expression of an HMG CoA
30 reductase in conjunction with DNA sequences for a steroid pathway enzyme is expected to result in alteration of the steroid pathway product profiles in oil as the enhanced steroid pathway throughput produces substrates for the enhanced enzyme activity. The novel

phytostanol ester compositions, e.g., sitostanol ester compositions, thus produced constitute another aspect of the present invention.

5 A. Harvest of Steroid Compounds

Methods for the derivation of steroid compounds from cells are well known in the art. The invention contemplates the recovery of biosynthesized steroid compounds from the leaves and/or stems of plants, from
10 plant seeds, from plant's vegetative organs, from callouses, and from cell cultures of plants, yeasts or eukaryotic cells.

Different sources of steroid compounds are preferred for various plants. For use as a food or a
15 food component as discussed later, the steroid compounds need not be isolated or purified to 100 percent purity. Steroid compound-enriched plants may be utilized directly.

For example, from tobacco or *Arabidopsis*, it may
20 be preferable to extract a pulp of the leaves and stems. From tomato, potato, or corn, it may be preferable to use the tomato, potato or corn in the form of the familiar storage organs that are typically consumed either directly, or a derivative thereof, such
25 as tomato paste, potato flakes, vegetable oil and many more that are well known in the food science arts.

If desired, after cultivation, the transgenic plant is harvested to recover the sterol or squalene product. This harvesting step can consist of
30 harvesting a callus culture, the entire plant, or only the leaves, or roots of the plant. This step can either kill the plant or, if only a non-essential portion of the transgenic plant is harvested, can permit the remainder of the plant to continue to grow.

100299" 22258350

Table 1. Demographic characteristics of the study population	
Characteristic	Frequency (%)
Age (years)	
< 18	10 (10.0)
18-24	15 (15.0)
25-34	20 (20.0)
35-44	25 (25.0)
45-54	30 (30.0)
55-64	35 (35.0)
65-74	40 (40.0)
75-84	45 (45.0)
85-94	50 (50.0)
≥ 95	55 (55.0)
Gender	
Male	60 (60.0)
Female	40 (40.0)
Ethnicity	
White	30 (30.0)
Black	20 (20.0)
Hispanic	10 (10.0)
Asian	5 (5.0)
Other	5 (5.0)
Marital status	
Married	40 (40.0)
Single	30 (30.0)
Divorced	20 (20.0)
Widowed	10 (10.0)
Education level	
High school or less	20 (20.0)
Some college	15 (15.0)
Bachelor's degree	25 (25.0)
Master's degree	10 (10.0)
Doctorate	5 (5.0)
Income (USD/year)	
< 10,000	10 (10.0)
10,000-19,999	15 (15.0)
20,000-29,999	20 (20.0)
30,000-39,999	25 (25.0)
40,000-49,999	30 (30.0)
50,000-59,999	35 (35.0)
60,000-69,999	40 (40.0)
70,000-79,999	45 (45.0)
80,000-89,999	50 (50.0)
90,000-99,999	55 (55.0)
≥ 100,000	60 (60.0)

- (iii) isolating the squalene or sterol(s) from the solution or suspension.

25 At least a portion of the transgenic plant is homogenized to produce a plant pulp using methods well known to one skilled in the art. This homogenization can be done manually, by a machine, or by a chemical
30 means as long as the transgenic plant portions are broken up into small pieces to produce a plant pulp. This plant pulp consists of a mixture of squalene or the sterol of interest, residual amounts of precursors, cellular particles and cytosol contents. This pulp can

be dried and compressed into pellets or tablets and eaten or otherwise used to derive the benefits, or the pulp can be subjected to extraction procedures.

The sterol or squalene can be extracted from the plant pulp produced above to form a sterol-or-squalene-containing solution or suspension. Such extraction processes are common and well known to one skilled in this art. For example, the extracting step can consist of soaking or immersing the plant pulp in a suitable solvent. This suitable solvent is capable of dissolving or suspending the squalene or sterol present in the plant pulp to produce a sterol-or squalene-containing solution or suspension. Solvents useful for such an extraction process are well known to those skilled in the art and included several organic solvents and combinations thereof such as methanol, ethanol, isopropanol, acetone, acetonitrile, tetrahydrofuran (THF), hexane, and chloroform as well as water-organic solvent mixtures. A vegetable oil such as peanut, corn, soybean and similar oils can also be used for this extraction as can steam distillation.

A whole plant or callus culture with an added, exogenous structural gene for a polypeptide having HMG-CoA reductase activity is grown under suitable conditions for a period of time sufficient for squalene or sterols to be synthesized and accumulated. The sterol-squalene-containing plant cells, preferably in dried form, are then lysed chemically or mechanically, and the squalene or sterol is extracted from the lysed cells using a liquid organic solvent or steam distillation, as described before, to form a sterol- or squalene-containing liquid solution or suspension by usual means such as chromatography.

09885722-062904

[illegible]

15 B. Harvest of Oil

The novel biosynthetic composition of the oil in the transgenic plants is contemplated. Thus, the present invention contemplates oil containing at least one sterol pathway product, extracted from seed of a described transgenic plant. Preferably, sitosterol, at least one sitosterol ester, or mixtures thereof, comprise at least about 50% by weight of the total sterol compounds of the oil. Preferably, sitosterol, at least one sitosterol ester, or mixtures thereof, comprise at least about 0.08% of the dry weight of a contemplated seed. Preferably, the oil has a reduced amount of squalene, cycloartenol, 24-methylene cycloartenol, obtusifoliol, stigmasta-7-enol, campesterol, or mixtures thereof, compared to oil from a corresponding transgenic plant that does not contain introduced DNA encoding a squalene epoxidase enzyme, a sterol methyl transferase I enzyme, a sterol C4-demethylase enzyme, a obtusifoliol C14 α -demethylase

enzyme, a sterol C5-desaturase enzyme, a sterol methyl transferase II enzyme, or mixture thereof, and that reduction is in the range of from about 10% to about 100%.

- 5 Oil is extracted from transgenic plant seeds of the present invention by method well known in the art. By way of example, oil can be extracted from plant seeds using extraction methods set forth above for harvesting sterols and squalene from transgenic plants.
- 10 Alternatively, oil can be extracted from transgenic plant seeds by usually used methods for obtaining seed oils such as by crushing the seeds to produce a pulp and then pressing the pulp to obtain oil. The pulp can also be extracted with appropriate solvents (e.g.
- 15 benzene) to obtain the oil. *Industrial Chemistry: A Manual for the Student and Manufacturer*, ed. By A. Rogers and A. B. Aubert, D. Van Nostrand Co., New York, pages 547-548 (1912).

20 C. Uses of Oil

- As discussed in the "Description of Related Art," phytosterols such as sitosterol are beneficial for lowering serum cholesterol (Ling et al. (1995) *Life Sciences* 57: 195-206) and preventing cardiac disease.
- 25 Tocopherols act as antioxidants, and play a major role in protecting cells from damage caused by free radicals (Halliwell (1997) *Nutrition Review* 55: 44-60). As the amount of sitosterol in conventional vegetable and bran oils is low relative to that of other sterol compounds,
- 30 the oils of the present invention are particularly useful for reducing the concentration of low density lipoprotein cholesterol in plasma.

Thus, further aspects of the present invention include the following:

Cholesterol-lowering compositions comprising the oils and sitostanol ester compositions disclosed herein. Such cholesterol-lowering compositions can take the form of, or be used in, foods, food products, processed foods, food ingredients, food additive compositions, or dietary supplements that contain oils and/or fats. Non-limiting examples include margarines; butters; shortenings; cooking oils; frying oils; dressings, such as salad dressings; spreads; mayonnaises; and vitamin/mineral supplements. Patent documents relating to such compositions include U.S. Patents 4,588,717 and 5,244,887, and PCT International Publication Nos. WO 96/38047, WO 97/42830, WO 98/06405, and WO 98/06714. Additional non-limiting examples include toppings; dairy products such as cheese and processed cheese; processed meat; pastas; sauces; cereals; desserts, including frozen and shelf-stable desserts; dips; chips; baked goods; pastries; cookies; snack bars; confections; chocolates; beverages; unextracted seed; and unextracted seed that has been ground, cracked, milled, rolled, extruded, pelleted, defatted, dehydrated, or otherwise processed, but which still contains the oils, etc., disclosed herein.

Food additive compositions of the present invention can be made by a method comprising obtaining oil containing a phytostanol or phytostanol ester selected from sitostanol, at least one sitostanol ester, or mixtures thereof, from cultured cells, or seeds of a plant, of the present invention, and evenly distributing the oil or desired phytostanol compound in finely divided form throughout the food product or food additive composition to which it is added by dissolution or by suspension in an emulsion. For example, the oil or phytostanol compound can be

dissolved in an edible solubilizing agent, or can be mixed with an edible solubilizing agent, an effective amount of a dispersant, and optionally, an effective amount of an antioxidant. Examples of useful edible
5 solubilizing agents include, but are not limited to, monoglycerides, diglycerides, triglycerides, vegetable oils, tocopherols, alcohols, polyols, or mixtures thereof. Examples of useful antioxidants include, but are not limited to, tocopherols, such as α -tocopherol, ascorbic acid, inexpensive synthetic antioxidants, and
10 mixtures thereof. Effective carriers for preparing emulsions or suspensions include water, alcohols, polyols, other edible compounds in which the oil or phytostanol compound is soluble or insoluble, and mixtures thereof. Examples of useful dispersants
15 include, but are not limited to, lecithin, other phospholipids, sodium lauryl sulfate, fatty acids, salts of fatty acids, fatty acid esters, other detergent-like molecules, and mixtures thereof. Alternatively, the food additive composition can be made by a method comprising obtaining oil containing at least one tocopherol, and a phytostanol or phytostanol ester selected from sitostanol, at least one sitostanol ester, and mixtures thereof, from cultured cells, or
20 seed of a plant, of the present invention, and mixing the oil with an edible solubilizing agent and an effective amount of a dispersant. Again, the edible solubilizing agent can include, but is not limited to, monoglycerides, diglycerides, triglycerides, vegetable
25 oils, tocopherols, alcohols, polyols, or mixtures thereof, and the dispersant can include, but is not limited to, lecithin, other phospholipids, sodium lauryl sulfate, fatty acids, salts of fatty acids,

30

T00290" E2 253360

fatty acid esters, other detergent-like molecules, and mixtures thereof.

The cholesterol-lowering compositions can also take the form of pharmaceutical compositions comprising a cholesterol-lowering effective amount of the oils or sitostanol ester compositions disclosed herein, along with a pharmaceutically acceptable carrier, excipient, or diluent. These pharmaceutical compositions can be in the form of a liquid or a solid. Liquids can be solutions or suspensions; solids can be in the form of a powder, a granule, a pill, a tablet, a gel, or an extrudate. U.S. Patent 5,270,041 relates to sterol-containing pharmaceutical compositions.

Any of the foregoing cholesterol-lowering compositions can be used alone or in combination in methods to lower the risk of developing an elevated plasma concentration of low density lipoprotein cholesterol, to lower the plasma concentration of low density lipoprotein cholesterol, or to treat or prevent an elevated plasma concentration of low density lipoprotein cholesterol. Such methods comprise orally administering to a human or animal subject an effective amount of cholesterol-lowering composition. What constitutes an effective amount of cholesterol-lowering composition can be determined empirically, and depends in part on a variety of factors, including the age, weight, sex, diet, general medical condition of the subject, and the severity of hypercholesterolemia. Subjects undergoing treatment with the cholesterol-lowering combinations disclosed herein can be monitored by routine measurement of serum cholesterol levels to determine the effectiveness of therapy. Continuous analysis of the data obtained in this way permits modification of the treatment regimen during therapy so

that optimal effective amounts of the cholesterol-lowering compositions of this invention are administered, and so that the duration of treatment can be determined as well. In this way, the treatment regimen/dosing schedule can be rationally modified over the course of treatment so as to achieve the lowest cholesterol-lowering effective amount of the present compositions which results in satisfactory anti-cholesterolemic effectiveness, and so that administration of these compositions is continued only so long as is necessary to successfully treat this condition. In general, an effective amount of a cholesterol-lowering composition of the present invention in the form of a phytostanol- or phytostanol ester-containing composition is in the range of from about 0.1 gm/day to about 4.5 gm/day. By way of example, a phytostanol ester composition, for example a sitostanol ester composition, can be administered in an amount in the range of from about 0.1 gm/day to about 4.5 gm/day, preferably from about 1 gm/day to about 4.5 gm/day, more preferably from about 2 gm/day to about 4.5 gm/day. A phytostanol composition, for example a sitostanol composition, can be administered in an amount in the range of from about 0.1 gm/day to about 3 gm/day, preferably from about 1 gm/day to about 3 gm/day, more preferably from about 2 gm/day to about 3 gm/day.

The cholesterol-lowering compositions of the present invention can be administered daily to patients in accordance with a number of different regimens. Fundamentally, these compositions should be administered in a cholesterol-lowering effective amount for a period of time effective to exert their anti-hypercholesterolemic preventing, reducing, or reversing

09885723-062004
T00290-22258860

action. Administration of the present cholesterol-lowering compositions should be continued until the hypercholesterolemic condition has been controlled or eliminated.

5 Another method encompassed by the present invention is that of achieving or improving effective absorption of sitostanol into a host, comprising producing at least one sitostanol ester by any of the methods disclosed herein, and administering this
10 sitostanol ester to a host, which can be a human or animal. The sitostanol ester can be administered by a route selected from oral route, parenteral route, or topical route. The dose, which can be administered daily, can be up to about 10 milligrams of the
15 sitostanol ester per kilogram of body weight. U.S. Patent 5,202,045 relates to the use of stanol fatty acid esters to reduce serum cholesterol.

Also envisioned are plants which in addition to having increased levels of phytosterols and
20 phytostanols due to the presence of constructs comprising sequences encoding 3-hydroxy-3-methylglutaryl-Coenzyme A reductase and at least one other sterol synthesis pathway enzyme, have increase levels of tocopherol due to the presence of constructs
25 comprising sequences allowing the overexpression of enzymes in the tocopherol biosynthetic pathway.

Tocopherol levels vary in different plants, tissues, and developmental stages, indicating a highly regulated biosynthetic pathway. The production of
30 homogentisic acid by *p*-hydroxyphenylpyruvate dioxygenase is likely to be a regulatory point for bulk flow through the pathway because of irreversible enzyme action and because homogentisic acid production is the first committed step in tocopherol biosynthesis (Norris

098857-02004
000230-0225850

et al., 1995, *Plant Cell* 7: 2139-2149). The other key regulatory step in tocopherol biosynthesis is the availability of the phytylpyrophosphate pool. Feeding studies (Furuya et al., 1987, *Phytochem.*, 26: 2741-
5 2747) in safflower callus culture demonstrated 1.8-fold and 18-fold increases in tocopherol synthesis by feeding homogentisate and phytol, respectively. In meadow rescue leaf, vitamin E increases in the initial phase of foliar senescence when phytol is cleaved off
10 from the chlorophylls and when free phytol is available (Peskier et al., 1989, *J. Plant Physiol.* 135: 428-432). These reports suggest tight coupling of tocopherol biosynthesis to the availability of homogentisic acid and phytol.

15 Transformation of plants with nucleic acid constructs that increase the biosynthetic activity of the tocopherol pathway can lead to increased production of particular tocopherol isomers, for example, α -tocopherol, are known in the art and can be found, for
20 example, in PCT International publication WO 00/61771 which is incorporated herein by reference. Formation of α -tocopherol from other tocopherols occurs due to S-adenosylmethionine (SAM)-dependent methylases such as γ -tocopherol methyl transferase. Overexpression of
25 methyl transferases in combination with 3-hydroxy-3-methylglutaryl-Coenzyme A reductase and at least one other sterol synthesis pathway enzyme as described herein is also contemplated in the present methods. Thus, any of the DNAs encoding enzymes of the
30 tocopherol biosynthetic pathway, discussed above, are useful in the present invention. Transformation of plants with an early tocopherol biosynthesis gene is sufficient to produce plants having an elevated level

T00290" E225850

of tocopherols. By "early tocopherol biosynthesis gene" is meant DNA encoding geranylgeranylpyrophosphate synthase, geranylgeranylpyrophosphate hydrogenase, 4-hydroxyphenylpyruvate dioxygenase, and phytyl/prenyl transferase. DNA encoding enzymes active in later steps of tocopherol biosynthesis ("secondary tocopherol biosynthesis genes") can be expressed to enhance carbon flux through the tocopherol pathway even further, and to produce specific tocopherol isomers. In this way, the tocopherol biosynthetic pathway can be modified to enhance production of any tocopherol compound of interest, such as α -tocopherol. As noted above, a variety of sources are available for the early tocopherol biosynthesis genes (and other tocopherol biosynthesis genes), and a gene from any of these sources can be utilized. If co-suppression occurs when a plant gene native to the target host plant is used to increase expression of a particular enzyme, a coding sequence from another source can be used as an alternative.

Preferred genes for introduction into plants to alter tocopherol quantity/quality include 3-deoxy-D-arabino-heptulosonate-7-P synthase (DAHP synthase), shikimate kinase, either or both of the prephenate dehydrogenases, 1-deoxy-d-xylulose 5-phosphate synthetase (DXS), 1-deoxy-d-xylulose 5-phosphate reductoisomerase (DXR), 4-diphosphocytidyl-2C-methyl-d-erythritol synthase (YgbP), 4-diphosphocytidyl-2C-methyl-d-erythritol kinase (YchB), 2C-methyl-d-erythritol 2,4-cyclodiphosphate synthase (YgbB), the gene product of GcpE, LytB (Altincicek et al., 2001, *J. Bacteriol.*, 183:2411-2416; Altincicek et al., 2001, *J. Immunol.*, 166:3655-3658; Campos et al., 2001, *FEBS*

Lett., 488:170-173), geranylgeranylpyrophosphate synthase (GGPPS), geranylgeranylpyrophosphate hydrogenase (GGH), phytyl/prenyltransferase (PPT), 4-hydroxy-phenylpyruvate dioxygenase (HPPD), 2-methyl-6-phytylplastoquinol tocopherol methyltransferase I (MTI), tocopherol cyclase, γ -tocopherol methyltransferase (GMT) a plant *slr 1736* gene (see Cyanobase <http://www.kazusa.or.jp/cyanbase>), a plant *slr 1737* gene (see Cyanobase <http://www.kazusa.or.jp/cyanbase>), an *ATPT2* gene (Smith et al., *Plant J.*, 11:83-92, 1977), and an *AANT1* gene (Saint Guily et al., *Plant Physiol.*, 100:1069-1071, 1992).

4-hydroxy-phenylpyruvate dioxygenase and geranylgeranylpyrophosphate hydrogenase will increase the homogentisate and phytol pools, respectively. Enzymes that control fluxes through pathways are well known to be regulated in higher organisms such as plants. Therefore, 4-hydroxyphenylpyruvate dioxygenase and geranylgeranylpyrophosphate hydrogenase genes of microbial origin which are not subject to regulation in plants, or those from higher organisms (plants, algae, fungi, etc.) that are deregulated, are especially attractive in this regard. A non-limiting example is the microbial enzyme 4-amino-4-deoxyprephenate dehydrogenase (TyrA from *Erwinia herbicola*) which can replace prephenate aminotransferase, arogenate dehydrogenase and aminotransferase. Overexpression of enzymes such as 3-deoxy-arabino-heptulosonate 7-P (DAHP) synthase, prephenate dehydrogenase, and shikimate kinase would lead to increases in the levels of homogentisate. DNA encoding any of the tocopherol biosynthetic enzymes discussed herein can be introduced alone or in various combinations to enhance tocopherol

quantity and/or alter tocopherol quality. When introduction of multiple enzymes is desirable, preferred combinations include, but are not limited to, 4-hydroxyphenylpyruvate dioxygenase (HPPD) plus
5 geranylgeranylpyro-phosphate hydrogenase (GGH), geranylgeranylpyrophosphate synthase (GGPP synthase) plus geranylgeranylpyrophosphate hydrogenase (GGH), 4-amino-4-deoxyphenate dehydrogenase (TyrA) plus phytylprenyltransferase (PPT), geranylgeranylpyro-
10 phosphate hydrogenase (GGH) plus phytylprenyltransferase (PPT), geranylgeranylpyrophosphate synthase (GGPP synthase) plus phytylprenyltransferase (PPT), 2-methyl-6-phytylplastoquinol tocopherol methyltransferase I (MTI)
15 plus phytylprenyltransferase (PPT), or 2-methyl-6-phytylplastoquinol tocopherol methyltransferase I (MTI), phytylprenyltransferase (PPT), 4-hydroxyphenylpyruvate dioxygenase (HPPD) and geranylgeranylpyrophosphate synthase (GGPP synthase).
20 Plants characterized by increase levels of sterol and tocopherol production can be produced by transforming plant cells or tissues genetically altered for increased sterol production by the methods described herein with additional nucleic acid
25 constructs encoding tocopherol biosynthetic enzymes. Introduction of constructs encoding tocopherol pathway enzymes can be accomplished using standard methods in molecular biology such as those described herein or those described in PCT International Publication WO
30 00/61771. Introduction of constructs encoding 3-hydroxy-3-methylglutaryl-Coenzyme A reductase, at least one other sterol synthesis pathway enzyme, and at least one tocopherol synthesis pathway enzyme can be accomplished in a single transformation or in a series

15

25

30

methods for obtaining oil containing phytosterols,

phytosterol esters, phytostanols, phytostanol esters, or mixtures thereof along with tocopherols employing extraction and processing procedures known in the art. Note, in this regard, Kochhar (1983) *Prog. Lipid Res.* 22: 161-188. Alternatively, seeds with increased levels of tocopherols and phytosterols, phytosterol esters, phytostanols, phytostanol esters, or mixtures thereof; or fruits and vegetables with increased levels of tocopherols and phytosterols, phytosterol esters, phytostanols, phytostanol esters, or mixtures thereof, can be used directly.

Tocopherols and phytosterols and/or phytostanols can then be obtained from deodorized distillates of oil seed extracts and in particular soybean oil extracts. Such deodorized distillates are expected to contain increased levels of both tocopherols and phytosterols and/or phytostanol extracts. Oil extracts from plants and seed of the present invention are particularly valuable in that they allow the production of high sterol/tocopherol oils in a single process thus resulting in reduced purification costs, processing time and waste stream. Methods for the isolation of tocopherols and sterols from plant oils are well known in the art and can be found, for example, in U.S. Patent Nos. 4,454,329; 5,097,012; 5,594,437; and 5,981,781.

EXAMPLES

The following examples are intended to provide illustrations of the application of the present invention. The following examples are not intended to completely define or otherwise limit the scope of the invention.

Example 1. Enhancement of Phytosterol content in seeds of transgenic plants by seed-specific overexpression of full-length HMG-CoA reductase (HMGR)

In order to examine the ability of HMGR overexpression for increasing sterol compound levels in seeds, the following experiment was performed in *Glycine max*. A full-length HMGR gene from rubber genomic DNA was expressed in developing *Glycine max* seeds using the 7S promoter. This was achieved by excising the rubber HMGR gene from the plasmid pHEV15 (Schaller et al., (1995) *Plant Physiol.*, 109: 761-770) using *EcoRI*. The 3.8 Kb fragment was inserted into the *EcoRI* site of pMON29920 (Fig. 3) such that the HMGR gene is flanked by the 7S promoter on the 5' end and the E9 3' terminator to create pMON43800 (Fig. 4). This was next digested with *SalI* and *NotI* to release a 7.7 Kb fragment that was then blunt-ended at the *SalI* end before ligating to pMON23616 (Fig. 5) that was first cut with *SmaI* and *NotI*. This created the pMON43818 binary vector that contains the rubber HMGR gene driven by 7S promoter and the NPTII gene selection marker driven by the NOS promoter and 3' NOS terminator. PMON43818 (Fig. 6) was used to transform *Agrobacterium tumefaciens* and transform *Glycine max* cotyledon explants as described in Example 2.

Seeds from 15 transgenic plants and one nontransgenic control plant were harvested at maturity. Sterol extraction and analysis on ten individual seeds per plant were performed as described in Example 2. Results are presented in Table 2.

Event	Campesterol	Stig- asterol	Sitosterol	Sito- tanol	Others (Pathway intermediates)	Total	Inter- mediate accumul ation
	ug/g	ug/g	ug/g	ug/g	ug/g	ug/g	(% of total sterol)
1	161.9	148.2	551.3	36.8	264.8	1163	22.8
2	241.6	287.9	1128.8	96.6	1489.8	3244.5	46
3	442.4	320.1	1876.6	117.3	1728.4	4484.8	38.5
4	311.2	345.6	1645.6	113.8	1307.5	3723.6	35
5	395.5	323.0	1592.1	83.1	933.8	3327.5	28
6	370.5	301.6	1735.8	97.2	990.5	3495.6	28.3
7	351.0	307.0	1457.3	101.1	885.3	3101.7	28.5
8	248	172.4	1270.1	74.3	428.8	2193.6	19.5
9	221.1	140.7	1149	76.7	652.6	2240.1	29.1
10	234.2	184.8	1306.8	64.1	669.4	2459.3	27.2
11	156.5	125.4	679.2	38.8	142.3	1142.2	12.4
12	311.2	242.9	1457.3	67	418.6	2497	16.7
13	165.4	135.4	1320.1	59.7	1645.8	3326.4	49.4
14	190.8	152	1121.3	51.4	1040.7	2556.2	40.7
15	182.9	157.4	1118.5	55.2	376.6	1890.6	20
16	197.9	151.7	946.6	61.7	225.3	1583.2	14.2

Table 2: Sterol profile of transgenic soybean plants expressing rubber HMGR gene driven by 7s

5 promoter. Event 1: control, events 2-16: 15 transgenic plants generated by 15 independent events using *Agrobacterium* mediated transformation.

Total sterols increased by 3.2- and 3.9- fold in the best performing plants (transgenic events 3 and 4).
 10 These two events also showed the highest increases of individual sterols. Campesterol increased by 2.7-fold, sitosterol by 3.4-fold, sitostanol by 3.2-fold and other sterols by 6.5-fold in event 3 while stigmasterol increased by 2.3-fold in event 4. The other sterols,
 15 which account for the highest increase in total sterols were pathway intermediates that included squalene, cycloartenol, 24-methylene cycloartenol, obtusifoliol, isofucosterol, and stigmasta-7-enol. These pathway intermediates normally form minor constituents in the

sterol composition of seeds. However, in the transgenic seeds, probably due to increased carbon flux through the pathway, they accumulate in significant amounts. This suggests additional control points for sterol biosynthesis in plants such as squalene epoxidase, C-24 sterol methyltransferase, and C-14 obtusifolliol demethylase.

Example 2: Enhance phytosterol biosynthesis in seeds of transgenic soybean plants by seed-specific expression of catalytic domain of HMG-CoA Reductase (HMGR) alone and in combination with sterol methyl transferase II (SMTII)

In another embodiment of the present invention, the levels of sterol compounds, including sitosterol, sitostanol, campesterol, stigmasterol and at least one ester for each of the sterol compounds and mixture thereof, can be elevated in plant seeds by overexpression of catalytic domain of plant-HMG-CoA reductases. One can transform a plant of interest using an expression cassette or vector comprising DNA encoding a polypeptide exhibiting 3-hydroxy-3-methylglutaryl-Coenzyme A reductase (HMG-CoA reductase or HMGR) activity. HMGR cDNAs from rubber have been successfully used to increase plant sterol levels in plant tissues (Schallet et al. (1995) *Plant Physiol.* 109: 761-770). Full-length and truncated forms of HMGR CDNAs encoding full-length and catalytic domain of HMGR, respectively, from *Arabidopsis* have also been used to overproduce sterols in transgenic *Arabidopsis* plants (Gonzalez et al. (1997) *Third Terpnet Meeting of the European Network on Plant Isoprenoids Abstracts*, Abstract No. 33, page 33). In the above examples however, the genes have not been specifically targeted

[illegible]

In order to examine whether overexpression of the catalytic domain of HMGR increases sterol levels in the seeds of transgenic soybean, the following experiment was performed in *Glycine max*. A truncated form of HMGR1 cDNA encoding only the catalytic domain of HMGR from *Arabidopsis* was expressed in developing seeds of *Glycine max* using the seed-specific 7S promoter. This was achieved by excising the cDNA fragment (HMGR1cd) encoding the HMGR1 catalytic domain from the plasmid pHMGR1cd (Dale et al., (1995) Eur. J. Biochem. 233: 506-513) using NdeI and SmaI enzymes resulting in the isolation of a 1.9 Kb fragment. The NdeI overhang was filled-in and the 1.9 Kb fragment was blunt-end ligated to vector pMON43818 (Figure 6), previously XhoI (XhoI overhang was filled-in) and SmaI digested such that the HMGR1cd was flanked by the 7S promoter on the 5' end and the E9 3' terminator to create a recombinant vector pMON43052 (Figure 7). This was next digested with XbaI and blunt-ended and then

digested with NotI to release a 3.4 Kb fragment and
ligated to pMON51850 (Figure 8) that was digested with
SmaI and NotI. The ligation created a recombinant
binary vector pMON43057 (Figure 9) that contained the
5 cDNA fragment encoding the catalytic domain of
Arabidopsis HMGR1, driven by 7S promoter and E9 3'
terminator and the NPTII selectable marker gene driven
by the NOS promoter and 3' NOS terminator. The
pMON43057 was used for *Agrobacterium tumefaciens*
10 mediated transformation of *Glycine max* cotyledon
explants. The pMON43058 (Figure 10) construct carrying
both the catalytic domain of *Arabidopsis* HMGR1 and
Arabidopsis SMTII, both driven by the 7S promoter, was
also used for *Agrobacterium tumefaciens*-mediated
15 transformation of *Glycine max* in a similar manner
described below.

Explants for transformation were prepared as
follows: sterilized seeds were germinated on
germination medium under light at 28°C for 5-6 days.
20 Germinated seeds were placed in the dark at 4°C for 24
hours prior to excision. Seed coats were removed and
hypocotyls of each seedling trimmed to a length of 0.5
cm to 1.0 cm in length. The cotyledons were then split
open such that the hypocotyl was split down in the
25 middle. The primary leaves and apical region of each
cotyledon was removed to expose the wounding region.
Wounding was performed with 3-7 shallow, scalpel scores
in line with the embryo axis, ensuring that the apical
bud was damaged. Wounded explants were incubated in
30 the culture of *Agrobacterium tumefaciens* containing
pMON43057. Incubation was for 1 hour at room
temperature. Inoculated explants were then
transferred to a co-culture medium and placed under
light at 23°C for 3-4 days. At this time explants were

T00290" E2458560

transferred to shooting medium without kanamycin selection and placed in a 25°C light growth room for 4 days.

After 4 days on delay, explants were transferred
5 to a 186 ppm kanamycin selection medium and placed in a 25°C light growth room for 2 weeks. At the end of two weeks explants were transferred to 186 ppm Woody Plant medium and placed again in a 25°C light growth room for another 2 weeks. Cultures were transferred every 2
10 weeks to fresh medium for approximately 18-21 weeks. At the 6 week transfer, the cotyledons and any dead material were removed from the explants, and the petiole was cut. At each subsequent 2 week transfer, the petiole was cut to expose fresh cells to the
15 medium.

Transgenic shoots that were approximately ½" in length, with 2 nodes, 1 open trifoliate and an active growing point were selected, cut and transferred to rooting medium. Once a good root system was developed,
20 the plants were sent to the greenhouse to grow up in soils in pots.

Seeds from the 15 transgenic plants and one nontransgenic control plant were harvested at maturity. Ten individual seeds from each plant were weighed and
25 ground into fine powder using an electric grinder. A known amount of cholestane (usually 100µg in 100µl ethanol) was added to each approximately 50mg powder sample. Sterol compounds were hydrolyzed directly from the ground tissue by saponification with 2ml of 10% KOH
30 in methanol by refluxing the material at 60°C for 30 minutes. The refluxed samples were cooled to room temperature and filtered through glass wool. An equal volume of water was added to each filtrate, and the

1002290 "E2458860

nonsaponifiables were extracted by partitioning three times with equal volumes of hexane. The hexane phases were pooled and evaporated. The residues were resuspended in 1 ml of acetone, and quantitatively transferred to glass GC vials that were immediately capped. Sterols were analyzed by Gas Chromatography-Flame Ionizing Detector using the following conditions: Inlet temperature of 220°C, detector temperature of 320°C, and column oven temperature programmed from 220°C to 320°C with initial temperature for 1 minute and final temperature for 16 minutes and ramp rate of 8°/min. The column used was a glass capillary DB-5 column of 50 m length, 320 µm diameter, and a film thickness of 0.25 µm. The carrier gas was helium at a flow rate of 1.0ml/min. Results are presented in Table 3 and Table 4.

TABLE 3

PlantSeed ID	Countdown	Squellene	Unknown 1	Camphorol	Sigmatol	Unknown 2	Octadecol	Slostol	Sigmatol	Isocamphorol	Sigmatol-7-endo	Unknown 3	Cyclohexanol	24 Methylene Cyclohexanol	Total steroid	Total End Product	Total Intermediates	Total Ethyl Steroids	Total Methyl Steroids	Ratio of methyl:ethyl	Total Unknowns	% Intermediates (of Total Steroids)
Control	Control	33.4	16.5	152.0	153.6	0.0	0.0	547.1	34.6	15.9	31.8	24.0	47.8	28.5	1088.9	922.1	125.4	736.5	119.5	0.06	389.7	11.5
OM_A12866	235.2	17.7	119.5	130.4	98.4	94.4	82.3	68.9	273.8	274.8	5770.0	2637.8	2637.8	3232.5	2103.0	2057.8	3232.5	1988.1	149.0	0.08	389.7	57.6
OM_A12867	205.9	13.9	146.0	129.2	95.9	11.9	97.6	111.9	70.3	116.6	900.4	273.8	108.4	42.9	5094.0	2016.5	2103.0	1988.1	149.0	0.08	389.7	55.6
OM_A12783	228.3	20.8	178.3	140.0	99.7	84.2	71.7	123.7	79.1	164.5	245.9	203.9	23.8	48.4	5531.1	2182.8	3236.6	2169.0	178.3	0.08	360.4	56.0
OM_A12868	181.3	19.1	200.4	170.3	85.3	71.3	65.3	117.7	95.1	125.3	492.3	83.5	233.2	48.4	5531.1	2478.9	2886.3	2400.1	200.4	0.08	167.9	52.2
OM_A12869	2431.0	13.5	135.5	153.9	78.3	75.8	108.2	42.6	140.3	161.3	509.0	121.3	321.6	441.0	5537.3	1891.3	3432.7	1899.1	153.5	0.08	213.3	57.6
OM_A12870	2484.6	2.2	146.7	120.8	44.3	61.9	159.0	42.6	120.3	549.0	186.2	205.2	411.2	5285.0	2009.1	3043.2	1892.6	146.7	0.07	232.7	57.6	
OM_A12879	2244.6	8.2	131.8	122.1	41.5	79.3	93.7	13.7	210.5	403.2	315.6	403.2	315.6	403.2	2879.9	1127.7	2879.9	1127.7	131.9	0.12	106.2	54.2
OM_A12888	2038.8	13.6	137.1	121.7	41.5	79.3	93.7	13.7	210.5	403.2	315.6	403.2	315.6	403.2	2879.9	1127.7	2879.9	1127.7	131.9	0.12	106.2	54.2
OM_A13241	1945.6	11.9	145.7	118.7	89.0	60.1	105.4	85.0	149.2	594.6	209.0	229.5	481.1	9185.0	2046.6	2456.6	2184.2	145.7	0.07	281.8	35.1	
OM_A13242	1870.4	21.3	149.7	118.7	89.0	60.1	105.4	85.0	149.2	594.6	209.0	229.5	481.1	9185.0	2046.6	2456.6	2184.2	145.7	0.07	281.8	35.1	
OM_A13285	2335.6	7.6	138.6	149.5	84.7	118.6	147.0	58.7	124.2	547.0	278.7	315.6	303.5	3202.2	2559.5	2895.8	2559.5	158.6	0.05	369.0	47.8	
OM_A13270	1328.8	27.6	142.8	149.5	84.7	118.6	147.0	58.7	124.2	547.0	278.7	315.6	303.5	3202.2	2559.5	2895.8	2559.5	158.6	0.05	369.0	47.8	
OM_A13241	950.0	7.4	178.8	130.7	41.8	35.6	80.7	34.4	88.1	407.8	46.8	107.3	326.3	303.5	3202.2	2559.5	2895.8	2559.5	158.6	0.05	369.0	47.8
OM_A13242	2437.9	0.0	158.3	130.7	41.8	35.6	80.7	34.4	88.1	407.8	46.8	107.3	326.3	303.5	3202.2	2559.5	2895.8	2559.5	158.6	0.05	369.0	47.8
OM_A13243	1873.7	17.9	162.3	133.0	71.2	21.6	105.2	42.2	145.2	1016.6	177.3	153.4	332.3	482.4	5078.7	2076.1	2787.8	1729.1	170.8	0.12	133.6	57.3
OM_A13244	1740.1	19.0	162.3	133.0	71.2	21.6	105.2	42.2	145.2	1016.6	177.3	153.4	332.3	482.4	5078.7	2076.1	2787.8	1729.1	170.8	0.12	133.6	57.3
OM_A13245	1924.4	20.8	162.3	133.0	71.2	21.6	105.2	42.2	145.2	1016.6	177.3	153.4	332.3	482.4	5078.7	2076.1	2787.8	1729.1	170.8	0.12	133.6	57.3
OM_A13250	1864.5	21.8	162.3	133.0	71.2	21.6	105.2	42.2	145.2	1016.6	177.3	153.4	332.3	482.4	5078.7	2076.1	2787.8	1729.1	170.8	0.12	133.6	57.3
OM_A13257	1944.5	23.8	162.3	132.4	34.9	36.9	122.2	70.2	132.7	501.5	128.9	199.5	416.8	4993.7	1916.9	2811.9	1907.2	163.3	0.07	217.2	55.5	
OM_A13258	2013.5	10.7	151.9	130.1	61.8	27.4	130.4	54.3	128.2	73.7	80.0	223.9	591.9	5517.3	2177.8	4035.7	2359.4	152.1	0.07	182.6	52.2	
OM_A13259	2753.8	18.1	135.1	128.6	40.2	75.1	111.5	77.4	190.3	642.2	201.8	264.4	750.3	6447.6	2313.9	4035.7	2359.4	152.1	0.07	182.6	52.2	
OM_A13264	2248.8	19.6	184.3	128.0	29.1	30.0	912.8	33.6	78.5	4628.8	1137.2	137.2	309.9	4628.8	1985.7	2802.5	1579.6	164.3	0.08	298.1	60.7	
OM_A13269	1161	17	173	136	40	31	911	42	85	356	151	151	314	5356	1650	1741	1562	173	0.14	150	49.2	
averageHigh expression	2020	15	180	142	69	54	1096	53	147	532	671	157	151	532	5356	2112	3003	2098	180	0	242	56.1

TABLE 4

Construct	PlantSeed ID	Squalene ug/g	Unknown 1 ug/g	Compound 1 ug/g	Signal 1 ug/g	Unknown 2 ug/g	Observed ug/g	Signal 2 ug/g	Signal 3 ug/g	Signal 4 ug/g	Isocolic acid ug/g	Stigmasta 7 and ug/g	Unknown 3 ug/g	Cyclohexyl ug/g	At Methylene Cyclohexyl ug/g	Total sterol ug/g	Total End Product	Total Intermediates	Total Ethyl Sterols	Total Methyl Sterols	Ratio of methyl/ethyl	Total Unknowns	% Intermediates	
Control	Control	81.0	12.5	164.6	174.1	0.0	0.0	529.8	11.2	2.7	61.3	67.5	22.5	5.3	10.5	43.4	1027.7	902.2	107.7	740.4	164.8	0.22	17.8	10.5
pMON43058	GM_A1282	2714.8	20.8	78.0	133.2	30.3	30.3	1149.8	70.3	61.3	67.5	22.5	193.5	216.1	467.3	5891.6	2106.7	3540.2	2090.0	78.0	0.04	244.6	60.1	
pMON43058	GM_A1282	862.6	16.6	163.0	169.0	9.9	9.9	1039.6	36.2	24.0	28.5	39.7	118.5	267.6	3052.2	1694.3	1291.7	1555.3	163.0	0.10	66.2	42.3		
pMON43058	GM_A1281	2314.7	15.0	51.4	125.3	7.1	47.5	988.3	59.4	70.4	928.8	248.5	192.4	388.3	5437.2	2153.2	3013.4	2172.2	51.4	0.02	270.6	55.4		
pMON43058	GM_A1241	1619.2	23.6	89.3	136.5	28.7	50.6	1033.1	69.1	61.8	674.9	167.3	186.0	560.5	4920.6	2002.9	2678.1	1975.4	89.3	0.05	239.5	54.4		
pMON43058	GM_A1241	1045.3	24.1	160.1	149.9	28.6	88.7	887.5	39.0	48.4	49.5	52.7	105.7	266.6	3331.8	1731.8	1494.7	1620.1	160.1	0.10	105.4	44.9		
pMON43058	GM_A1242	1099.2	21.9	103.1	159.2	34.8	69.0	1098.9	63.2	78.9	85.1	173.1	212.0	786.0	5657.9	2273.2	3155.0	2249.0	103.1	0.05	229.7	55.8		
pMON43058	GM_A1242	1055.4	13.2	92.9	144.0	21.4	86.8	1198.0	92.2	59.5	497.7	196.6	155.0	730.9	4343.7	2024.8	2087.8	1991.4	92.9	0.05	231.1	48.1		
pMON43058	GM_A1242	2179.8	29.4	197.9	185.8	81.2	60.1	1286.6	75.5	153.8	485.4	191.1	323.3	752.4	5500.8	1855.0	3344.2	1810.7	197.9	0.11	301.6	60.8		
pMON43058	GM_A1242	1800.1	31.2	149.9	172.4	104.0	90.5	1425.2	84.6	280.0	826.4	213.2	191.1	783.2	5630.5	2530.0	3301.5	2660.1	149.9	0.06	349.0	53.4		
pMON43058	GM_A1242	1895.8	24.7	87.6	154.3	27.7	79.9	1425.2	84.6	75.1	788.0	213.2	191.1	783.2	5630.5	2539.7	2825.2	2527.3	87.6	0.03	265.7	50.2		
pMON43058	GM_A1242	2089.5	20.6	77.5	157.6	39.0	77.8	1350.2	68.8	88.8	95.6	172.7	265.7	827.5	6193.4	2011.7	3349.3	2623.0	77.5	0.03	232.4	54.1		
pMON43058	GM_A1242	2051.3	30.4	74.3	151.7	19.5	68.8	1409.4	57.2	65.0	88.3	137.6	161.8	876.9	6081.2	2682.0	3221.7	2672.7	74.3	0.03	177.5	51.7		
pMON43058	GM_A1242	2004.7	30.1	123.5	174.9	25.8	65.4	1347.9	50.9	73.4	1029.9	101.5	187.3	761.0	5976.4	2682.0	3061.8	2687.0	123.5	0.05	157.5	51.7		
pMON43058	GM_A1242	2047.7	24.1	102.6	178.5	32.7	85.3	1619.4	70.4	77.3	800.1	131.8	181.7	711.3	5930.6	2771.0	2970.4	2745.7	102.6	0.04	189.1	50.1		
pMON43058	GM_A1244	1914.8	24.6	102.6	178.5	32.7	85.3	1619.4	70.4	77.3	800.1	131.8	181.7	711.3	5930.6	2771.0	2970.4	2745.7	102.6	0.04	189.1	50.1		
pMON43058	GM_A1244	902.2	21.1	151.1	151.0	8.2	17.6	984.6	61.9	16.6	199.1	63.6	102.2	264.6	2943.7	1547.7	1303.2	1413.1	151.1	0.11	92.8	44.3		
pMON43058	GM_A1242	2392.3	22.5	86.6	145.8	7.9	82.7	1290.4	63.3	58.6	839.7	173.2	155.0	692.5	5890.6	2425.8	3361.1	2397.9	86.6	0.04	203.6	56.1		
pMON43058	GM_A1242	1808.9	22.5	111.8	155.8	31.7	81.9	1188.1	66.1	80.8	709.3	155.6	189.5	610.9	5101.4	2220.8	2751.8	2198.8	111.8	0.1	209.8	52.2		
43058 ever 10x																								

Hex

of the 10 transgenic lines with the pMON43058 showed 3- to 5-fold increase in total sterols. The best performing transgenic lines showed about 2- to 3-fold increase in sitosterol and 4.5- to 6-fold increase in sitostanol levels. However, the campesterol accumulation was reduced by 50% in these lines. This was due to overexpression of the *Arabidopsis* SMTII enzyme which enhances the carbon flux towards the synthesis of 24-ethyl sterols thereby reducing the carbon flux through the pathway leading to the synthesis of 24-methyl sterols. As seen in pMON43057 transgenic lines, all of the transgenic lines harboring the pMON43058 also accumulated 50-60% of the total sterols in the form of pathway intermediates which are squalene, cycloartenol, 24-methylene cycloartenol, obtusifoliol, isofucosterol, and stigmasta-7-enol. These pathway intermediates normally form minor constituents in the sterol composition of seeds. However, in the transgenic seeds, probably due to increased carbon flux through the pathway, they accumulate in significant amounts. The pathway intermediates accumulation is highly significant when the truncated form of HMGR is overexpressed as compared to the full length form of HMGR suggesting that the overexpression of the truncated form of HMGR creates even greater increase in carbon flux through the pathway. This provides further evidence for additional control points for sterol biosynthesis in plants such as squalene epoxidase, sterol methyltransferase I, sterol C4-demethylase, obtusifoliol C14 α -demethylase, sterol C5-desaturase, and sterol methyl transferase II.

190290 22 25850

Example 3. Enhancement of phytosterol biosynthesis in seeds of *Arabidopsis* transgenic plants by constitutive expression of different forms of *Arabidopsis* and rubber HMGR enzymes.

5

Arabidopsis transgenic plants were generated using *Agrobacterium* mediated transformation of constructs (pMON53733, pMON53734, pMON53735, pMON53736, pMON53737, pMON53738, pMON53739, pMON53740) carrying cDNA encoding different forms of *Arabidopsis* and rubber HMGR enzymes driven by CaMV enhanced 35S promoter (Figures 13-20). The transformed *Arabidopsis* seeds carrying each of the above constructs were selected on kanamycin (50µg/ml) medium to select for transformants expressing the

10

selectable marker, the NPTII gene driven by CaMV 35S promoter. Kanamycin resistant *Arabidopsis* transgenic plants were grown in green house for maturity and seeds were collected from each of the transgenic lines for sterol analysis. About 50 mg of seeds from each

15

transgenic line were weighed, homogenized and used for saponification to extract sterols as described in Example 2.

20

Figures 21-26 describe the sterol analysis data obtained from the transgenic lines carrying each of the above constructs. Figure 27 shows the effect on different sterol end products and pathway intermediate accumulation when different forms of rubber HMGR cDNAs were expressed constitutively in transgenic *Arabidopsis*

25

plants. When truncated rubber HMGR (with or without linker region) was overexpressed the total sterol accumulation in seeds increased by 2.9 to 3.7-fold as compared with the wild type control plants. The sterol end products such as campesterol and sitosterol showed

30

09805723-052001

1.5 to 2 -fold increase in the lines expressing truncated form of rubber HMGR (with and without linker). However the sitostanol end product accumulation in the transgenic lines harboring the truncated form of rubber HMGR (with and without linker) was enhanced by 2.8 to 7-fold. There is a significant accumulation of pathway intermediates such as cycloartenol and 24-methylene cycloartenol in the seeds of the transgenic lines transformed with the truncated form of rubber HMGR (with and without linker region). The wild type control plants used in the experiment do not accumulate both of the pathway intermediates.

Example 4. Comparison of Steroid Compounds from HMGR Constructs in a Yeast HMGR1 Knockout Mutant

The effects on the sterol levels of the expression of various HMGR constructs expressed in a yeast HMGR1 knockout mutant were compared. Constructs containing a nucleic acid encoding the full length HMGR polypeptides from *Arabidopsis* and rubber were compared to those encoding a truncated *Arabidopsis* or rubber HMGR polypeptide that were lacking both the membrane binding and linker region of HMGR. The control yeast cells were transformed with a similar construct lacking a polypeptide encoding any form of HMGR.

Yeast cells transformed with *Arabidopsis* HMGR and rubber HMGR constructs accumulated approximately the same amounts of zymosterol and ergosterol, but more squalene than the control yeast.

Transformed yeast cells having rubber HMGR constructs accumulated about the same amount of ergosterol, but about twice as much squalene and zymosterol than the control yeast.

Transformed yeast cells having *Arabidopsis* tHMGR constructs accumulated three times as much squalene, twice as much zymosterol, and about 30 percent more ergosterol than the control yeast.

5 Transformed yeast cells having rubber tHMGR constructs accumulated three times as much squalene, four times as much zymosterol, and about 50 percent more ergosterol than the control yeast.

The data are shown in a Figure 28, "Plant HMGR1
10 Constructs in Yeast HMGR1 Knockout Mutant".

Example 5. Gene sequences for all genes listed in the application

The sequences obtained from the NCBI public
15 database are SEQ ID NO.: 1,2,3,20,21,22,23. These sequences are included in the appendix and denoted as follows:

Appendix A= SEQ ID NO. 1,
Appendix B= SEQ ID NO. 2,
20 Appendix C= SEQ ID NO. 3,
Appendix D= SEQ ID NO. 20,
Appendix E= SEQ ID NO. 21,
Appendix F= SEQ ID NO. 22,
Appendix G= SEQ ID NO. 23.

25

SEQ ID 1 = *Arabidopsis* squalene epoxidase protein sequence (Accession NO: AC004786) **See Appendix A**

SEQ ID 2 = *Arabidopsis* squalene epoxidase (Accession NO: N64916) **See Appendix B**

30 SEQ ID 3 = *Arabidopsis* squalene epoxidase (Accession NO: T44667) **See Appendix C**

SEQ ID 4 = *Arabidopsis* squalene epoxidase (clone ID: ATA506263) nucleotide sequence
35

09886723-062001
T002500"E258569

GAATTCCTCCGGGTCGACCCACGCGTCCGCTTATAGATAAGGATATGGCCTT
TACGAACGTTTGCCTATGGACGCTACTCGCCTTCATGCTGACTTGGACAGTGTTCT
TACGTACAAACAGGGGGAAGAAGGCGACGAGTTGGCGGATGCGGTGGTTGAAG
AGCGAGAAGACGGTGCTACTGACGTTATCATCGTTGGGGCTGGAGTAGGCGGCTC
5 GGCTCTCGCATATGCTCTTGCTAAGGACGGGCGTGGAGTCCATGTAATAGAGAGG
GACCTGAGAGAACCAGAGAGAATCATGGGTGAGTTTATGCAACCAGGAGGACGAC
TCATGCTCTCTAAGCTTGGTCTTGAAGATTGTTTGGAGGGAATAGATGCCCAAAA
AGCCACGGGCATGACAGTTTATAAGGACGGAAAAGAAGCAGTCGCATCTTTTCCC
GTGGACAACAACAATTTTCTTTTGTATCCTTCGGCTCGATCTTTTACAATGGCC
10 GATTCGTCCAACGATTGCGGCAAAAGGCTTCTTCTCTTCCCAATGTGCGCCTGGA
AGAAGGAACGGTGAAGTCTTTGATAGAAGAAAAAGGAGTGATCAAAGGAGTGACA
TACAAAAATAGCGCAGGCGAAGAAACAACAGCCTTGGCACCTCTCACTGTAGTAT
GCGACGGTTGCTACTCAAACCTTCGCCGGTCTCTTAATGACAACAATGCGGAGGT
TCTGTCATACCAAGTTGGTTTTATCTCAAAGAACTGTCAGCTTGAAGAACCCGAA
15 AAGTTAAAGTTGATAATGTCTAAACCTCCTTCACCATGTTGTATCAAATCAGCA
GCACCGACGTTTCGTTGTGTTTTGAAGTTCTCCCAACAACATTCTCTTATTTT
AAATGGTGAAATGGCTACTTTCGTGAAGAACAATATTGCTCCTCAGGTACCTTTA
AAACTCCGCAAAATATTTTGAAGGGATTGATGAAGGAGAACATATAAAGCCA
TGCCAACAAAGAAGATGACAGCTACTTTGAGCGAGAAGAAAGGAGTGATTTTATT
20 GGGAGATGCATTCAACATGCGTCATCCAGCAATCGCATCTGGAATGATGGTTTTA
TTATCTGACATTCTCAATTTACGCCGTCTTCTCCAGCCATTAAGCAACCTTGGCA
ATGCGCAAAAAATCTCACAAGTTATCAAGTCCTTTTATGATATCCGCAAGCCAAT
GTCAGCGACAGTTAACACGTTAGGAAATGCATTCTCTCAAGTGCTAGTTGCATCG
ACGGACGAAGCAAAAGAGGCAATGAGACAAGGTTGCTATGATTACCTCTCTAGTG
25 GTGGGTTTTCGCACGTCAGGGATGATGGCTTTGCTAGGCGGCATGAACCCTCGTCC
GATCTCTCTCATCTATCATCTATGTGCTATCACTCTATCCTCCATTGGCCATCTA
CTCTCTCCATTTCCCTCTCCCCTTGGCATTGTCATAGCCTTCGACTTTTTTGGTT
TGGCTATGAAAATGTTGGTTCCCATCTCAAGGCTGAAGGAGTTAGCCAAATGTT
GTTTCCAGTCAACGCCGCCGCGTATAGCAAAAGCTATATGGCTGCAACGGCTCTT
30 TAAACACTGGTGCTTTAAACTGCAAAATATAACACATATATAAATCCCGAATCT
TTGTGATTCTGCATATATTGTGTTCTACAATTATTCTCATATAAATGAAAATTGT
TCTACGTAAAAGTAAAAGAAGGAATTGTAATACTAATAAAACGAGTTTTTAATT
CTGTTGAATGCTTGTGTATATTGGTGAAAAAAGGAGGAGGAGGAGGAGGAGGAGG
AAAAAAGGGCGGCCGC

35

SEQ ID 5 = *Arabidopsis* squalene epoxidase (clone
ID: ATA506263) amino acid translation

EFFGRPTRPLIDKDMAFTNVCLWTLAFMLTWTVFYVTNRGKKATQLADA
40 VVEEREDGATDVIIVGAGVGGALAYALAKDGRRVHVIERDLREPERIMGEFMP
GGRLMLSKLGLDCLEGIDAQKATGMTVYKDGKEAVASFPVDNNNFPDPSARSF
HNRFVQRLRQKASSLPNVRLEEGTVKSLIEEKGVIKGVITYKNSAGEETALAPL
TVVCDGCYSNLRRSLNDNNAEVL SYQVGFISKNCQLEEPKLLKIMSKPSFTMLY
QISSTDVRCVFVLPNNIPSI SNGEMATFVKNTIAPQVPLKLRKIFLKGIDEGEH
45 IKAMPTKKMTATLSEKKGVILLGDAFNMRHPAIASGMMVLLSDILILRRLQLPLS
NLGNAQKISQVIKSFYDIRKPM SATVNTLGNAFSQVLVASTDEAKEAMRQGCYDY
LSSGGFR TSGMMALLGGMNPRPISLIYHLCAITLSSIGHLLSPFPSPGLIWHSLR
LFGLAMKMLVPHLKAEGVSQMLFPVNAAAYSKSYMAATAL*

50

SEQ ID 6 = *Arabidopsis* squalene epoxidase (clone

0985723.062001
T00290" E2758860

[illegible]

45 MAFTHVCLWTLVAFVLTWTVFYLTNMKKKATDLADTVAEDQKDGAADV I IVGAGV
GGSALAYALAKDGRRVHVIERDMREPERMMGEFMQPGGRLMLS KLGLQDCLEDID
AQKATGLAVYKDGKEADAPFPVDNNNFSYEPSARSFHNGRFVQQLRRKAFLSLNV
RLEEGTVKSLLEEKGVVKGVTYKNKEGEETTALAPLTVVCDGCYSNLRRSLNDDN
NAEIMSYIVGYISKNCRLLEEPEKLHLILSKPSFTMVYQISS TDVRCGFVLPENF
50 PSIANGEMSTFMKNTIVPOVPPKLRLKIFLKGIDEGAHIKVVP AKRMTSTLSK KKG

VIVLGDAFNMRHPVVASGMMVLLSDILILRRLQLPLSNLGDANKVSEVINSFYDI
RKPM SATVNTLGNAFSQVLIGSTDEAKEAMRQGVYDYLCSSGGFRTSGMMALLGGM
NPRPLSLVYHLCAITLSSIGQLLSPFPSPLRIWHSCLKLFGLAMKMLVPNLKAEGV
SQ MLFPANAAAYHKSYMAATT*

5

SEQ ID 8 = *Arabidopsis* squalene epoxidase (clone
ID: ATA102071) nucleotide sequence

AAATCATATTGAGAACAAATAGATTTGGTTATATATGGCTTTTACGCACG
10 TTTGTTTATGGACGTTAGTCGCCTTCGTGCTGACGTGGACGGTGTCTACCTTAC
CAACATGAAGAAGAAGGCGACGGATTTGGCTGATACGGTGGCTGAGGATCAAAAA
GACGGTGCTGCTGACGTCATTATCGTCGGGGCTGGTGTAGGTGGTTCGGCTCTCG
CATATGCTCTGCTAAGTGTGCGCCTGGAAGAAGGAACGGTGAAGTCTTTACTAGA
AGAAAAAGGAGTGGTCAAAGGAGTGACATACAAGAATAAAGAATGCCAACAAACA
15 ACAGCCTTGGCACCTCTCACTGTGGTATGCGACGGTTGCTAATCAAACCTTCGTC
GGTCTCTTAATG

SEQ ID 9 = *Arabidopsis* squalene epoxidase (clone
ID: ATA102071) amino acid translation

20

MAFTHVCLWTLVAFVLTWTVFYLTNMKKKATDLADTVAEDQKDGAADVII
VGAGVGG SALAYALLSVRLEEGTVKSLLEEKGVVKGVTYKNKECEQTTALAPLTV
VCDGC

25

SEQ ID 10 = *Arabidopsis* squalene epoxidase (clone
ID: ATA504158) nucleotide sequence

CACAAAGCAAAAAAATCTCTGTAAAAGCAGAACGATAATGGAGTCACAAT
30 TATGGAATTGGATCTTACCTCTTTTGATCTCTTCTCTCCTCATCTCCTTCGTCGC
TTTCTATGGATTCTTCGTCAAACCGAAGCGGAACGGTCTCCGTCACGATCGGAAA
ACTGTTTCTACCGTCACCTCCGACGTCGGATCTGTTAATATTACCGGAGATACTG
TCGCTGATGTCATTGTTGTTGGAGCTGGTGTGCTGGTTCTGCTCTTGCTTATAC
TCTTGGAAGGGGAAATTTAAACGCCGAGTTCATGTGATTGAAAGAGATTTATCG
35 GAGCCTGATCGTATTGTTGGGAGTTGTTACAGCCTGNGGGTTACCTCAAGTTAC
TGGAGTGTGGAATTGGAGATTGTGTGGAAGAAATAGATGCTCAGCNTGTGTATGG
TTATGCACTTTTTAAAAATGGG

SEQ ID 11 = *Arabidopsis* squalene epoxidase (clone
ID: ATA504158) amino acid translation

40

TKQKNLCKSRTIMESQLWNWILPLLISLLISFVAFYGFVVKPKRNLRLH
DRKTVSTVTSVGSVNITGDTVADVIVVGAGVAGSALAYTLGKGKFKRRVHVIER
DLSEPDRIVGELLQPXGYLKLLECGIGDCVEEIDAQXVYGYALFKNG

45

SEQ ID 12 = *Arabidopsis* obtusifolius C14 α -
demethylase nucleotide sequence (Accession NO:
complement, join AC002329:37461...38456,

T00290" 2258860

AC002329:39121...39546) (homolog of sorghum obtusifolios
C14 α -demethylase) nucleotide sequence

CGTGTTTTACAAATTTCTTTGTTGGTTTTCCACAGATTTAAAGAACCCT
5 AACGAGAGAAAAAATGGACTGGGATTACTATACGCTGTTGAAGACGAGTGTGGC
TATTATTATAGTGTGTTGTGGCCAACTCATAACCTCCTCCAAATCCAAGAAG
AAAACAAGTGTCGTCCCACTCCCTCCAGTTCCTCAAGCGTGGCCTCCATTTATCG
GATCCCTAATCCGCTTCATGAAAGGTCCAATAGTGCTACTTAGAGAGGAATATCC
10 TAAGCTTGGAAGTGTGTTTACAGTGAAGCTTCTTCAAAAAACATCACTTTTCTC
ATCGGTCCCGAAGTCTCGTCCCACTTTTTCACGCTTATGAATCTGAACTCAGCC
AGAAAGAAATTTACAAATTTAATGTGCCTACTTTTGGCCCCGAGTTGTGTTTGA
TGTTGACTATCCCGTTCGGATGGAGCAGTTCGGATTCTTCTCCAGCGCTCTCAAG
GATTACTTCTCAAAATGGGGAGAAAGTGGGGAAGTGGATCTAAAGGCCGAGTTAG
AGCGTCTAATCACCTTGACTGCTAGTAGATGTCTATTGGGTCGAGAAGTCCGTGA
15 CCAACTTTTGTATGATGTTGCTCCATTGTTCCATGACCTTGATAAAGGCATGCAA
CCCATAAGTGTCATCTTCCCAAAGCTCCCCATTCAGCTCACAATTGTCGTGACC
GTGCTCGCGGAAAGATTGCAAAAATCTTTTCAAACATCATAGCAACAAGAAAACG
CTCTGGTGACAAATCAGAGAACGACATGCTACAATGTTTCATCGACTCAAAGTAC
AAAGACGGTAGAGAGACAACTGAATCTGAAGTAACTGGTTTGCTCATTGCTGGTT
20 TGTTTGAGGACAACATACAAGCTCTATCACTGCCACATGGACCGGTGCTTATCT
AATTCAAAACAAACACTGGTGGTCCGCGGCTTTGGACGAGCAGAAGAACTGATT
GGAAAACATGGGGACAAGATCGACTACGATGTTTTGTCTGAGATGGATTTTCTGT
TTCGAGTGCAAAAGAAGCTTTAAGGCTTACCCTCAAAGATCTTACTGCTGAG
AACAGTACACAGTGATTTACCGTGACAACCTCGAGAAGGAAAGCAATATGAGATA
25 CCAAAGGGTCATATCGTTGCAACTTCTCCTGCATTGCGCAACCGCTTACCTCATG
TCTACAAAGATCCGGAATTTTATGATCCGGATAGATTTTCAAAGGAAAGAGAAGA
GGATAAAGCAGCTGGTTCGTGTTTATACATCTCTTGGGAGCTGGTAGGCACGAG
TGTCCTGGTGGATCATTTGCGTTCTTGACAGATCAAAGCCGTATGGTGTCACTTAT
TGAGAACTTTGAGCTTGAGTTAGTGTACCTTTCCCTGAAATCAACTGGAATGC
30 TTTGGTTCGTTGGTGCTAAAGGAAATGTCATGGTTCGTTACAAGCGTCGTCCCTTT
TCTTAA

SEQ ID 13 = *Arabidopsis* obtusifolios C14 α -
35 demethylase nucleotide sequence (Accession NO:
complement, join AC002329:37461...38456,
AC002329:39121...39546) (homolog of sorghum obtusifolios
C14 α -demethylase) amino acid translation

MDWDYYTLLKTSVAIIIVFVAKLITSSKSKKKTSVVPLPPVLQAWPPFI
40 GSLIRFMKGPIVLLREEYPKLGSVFTVKLLHKNITFLIGPEVSSHFFNAYESEL
QKEIYKFNVPFTFGPGVVDVDYVPRMEQFRFFSSALKDYFSKWGESGEVDLKAEL
ERLITLTASRCLLGREVRDQLFDDVAPLFHDLDDKGMQPISVIFPKLPPIAHNCRD
RARGKIAKIFSNIIATRKRSGDKSENDMLQCFIDSKYKDGRETTESEVTGLLIAG
45 LFAGQHTSSITATWTGAYLIQNKHWSAALDEQKKLIGKHGDKIDYDVLSEMDFL
FRSAKEALRLHPPKILLRTHVSDFTVTTREGKQYEIPKGHIVATSPAFAANRLPH
VYKDPENFDPRFSKEREDKAAGSCSYISLGAGRHECPGGSFAFLQIKAVWCHL
LRNFELELVSPFPEINWNALVVGAKGNVMVRYKRRPFS*

50 SEQ ID 14 = *Arabidopsis* obtusifolios C14 α -

0985723-062001

demethylase(clone ID: ATA101105) nucleotide sequence

5 GACACTATAGAAGAGCTATGACGTCGCATGCACGCGTACGTAAGCTCGGA
ATTCCGGCTCGAGCTTGTTCACAAAAAGATTACTTTTCTTATTGGTCCTGAAGTCT
CTGCTCATTFTTTTCAAAGCTTCTGAATCTGATCTTAGTCAGCAGGAAGTGTATCA
GTTCAATGTCCCTACTTTTGGTCCTGGAGTTGTTTTCGATGTTGATTATTCTGTT
TCGTCAGGAGCAGTTCGGTTCTTCACTGAGGCACTTAGAGTTAACAAGTTGAAGG
GTTATGTGGATATGATGGTTACTGAAGCTGAGGATTACTTCTCTAAATGGGGAGA
GAGTGGTGAAGTTGATATTAAGGTTGAGCTAGAGAGGCTCATCATCTTGACTGCA
10 AGTGATGTTTACTGGGTCGAGAAGTTCGTGATCAGCTTTTGTATGATGTCTCTGC
TTTGTTCATGACCTTGACAATGGAATGCTTCCCATCAGTGCTTCCCATCAGTGT
TCTCTTCCCATATCTCCCAATTCCAGCTCACCG

15 SEQ ID 15 = *Arabidopsis obtusifoliol* C14 α -
demethylase(clone ID: ATA101105) amino acid translation

HYRRAMTSHARVRKLGIRLELVHKKITFLIGPEVSAHFFKASESDLSQQE
VYQFNVPTFGPGVVFVDVYSVRQEQFGSSLRHLELTS

20 SEQ ID 16 = *Arabidopsis obtusifoliol* C14 α -
demethylase(clone ID: ATA202967) nucleotide sequence

TCGACCCCGCGTCCGCGGACGCGTGGGATCAGCTTCAAGCTTAAGAGAGC
TTCGAAAGCGAAAGCGACGATTTCTTCTCCATCGTGAGAGCAAATCTCCAGAGCC
25 GTTTTCTCTTCTTCTTCTTCTCCTCCTCGCGCCGTCTCTGAAACTCCATCATCGTAT
CAATCAAATTGCTTCTCCTCCAAATTGAAAAACAATGGAATTGGATTCCGAGAA
CAAATTGTTGAAGACGGGTTTGGTTATAGTGGCGACACTTGTTATAGCCAACTC
ATCTTCTCTTTCTTCACTTCTGATTCTAAGAAGAAGCGTCTTCTCCTACTCTTA
AAGCTTGGCCTCCATTGGTTGGAAGTCTTATCAAATCTTGAAAGGACCTATTAT
30 TATGCTTAGAGAGGAATACCCTAAGCTTGGAAGTGTGTTTACTGTTAATCTTGTT
CACAAAAGATTACTTTTCTTATTGGTCTGAAGTCTCTGCTCATTTTTCAAAG
CTTCTGAATCTGATCTTAGTCAGCAGGAAGTGTATCAGTTCAATGTCCCTACTTT
TGGTCTCGGAGTTGTTTTCGATGTTGATTATTCTGTTTCGTCAGGAGCAGTTTCGG
TTCTTCACTGAGGCACTTAGAGTTAACAAGTTGAAGGGTTATGTGGATATGATGG
35 TTAAGGTTGAGCTAGAGAGGCTCATCATCTTGACTGCAAGTAGATGTTTACTGGGT
CGAGAAGTTTCGTGATCAGCTTTTGTATGATGTCTCTGCTTTGTTCCATGACCTTG
ACAATGGAATGCTTCCCATCAGTGTTCTCTTCCCATATCTCCCAATTCCAGCTCA
CCGCCGTGCTGACCGTGCCCGAGAAAAGCTTTTCGGAGATTTTCGAAAAATCATT
40 GGGTCGAGAAAACGCTCTGGA AAAACAGAGAACGACATGCTGCAGTGTTTCATCG
AATCAAAGTACAAAGATGGTAGACAGACAACCGAATCTGAAGTCACTGGTTTGCT
CATTGCTGCTCTGTtTGCAGGACAACACACGAGCTCTATCACTTCCACCTGGACC
GGTGCTTATCTGATGCGATACAAAGAGTACTTCTCAGCTGCTCTTGATGAGCAGA
AGAACCTGATTGCGAAACATGGAGACAAGATCGATCATGATATCTTATCCGAGAT
45 GGATGTTCTCTACCGCTGCATTAAGGAAGCGTTGAGGCTTCACCCTCCACTCATC
ATGTTAATGAGAGCCTCGCACAGTGATTTTCAGCGTGACAGCTCGGGATGGAAAAA
CTTACGATATCCCAAAGGGTCACATCGTTGCAACCTCCCCTGCATTTGCCAACCG
CTTACCGCACATCTTCAAAGACCCCGACACCTACGACCCAGAAAGATTCTCCCCT
GGAAGAGAAGAGGACAAAGCCGCAGGGGCATTCTCGTACATTGCATTTCGAGGGG
50 GAAGGCACGGGTGCCTTGGAGAGCCGTTTGCTTACCTGCAGATCAAAGCCATATG

0985723.062004

GAGTCATTTGTTGAGGAACTTCGAGCTTGAGCTAGTTTCACCGTTCCTGAGATT
 GACTGGAACGCTATGGTGGTTGGAGTTAAAGGCAATGTGATGGTGCGTTACAAGA
 GGCgcCAGCTTCTTAAAGACAAGTTTAAAGGTTATTGCAGCTTTGGATTTTTCTC
 TCTGGTTTCTGCTTTGCTTTTGTCCCTCTCTGGTTTTAGTTTTGTTGTTGAATAA
 5 TTCTTCTGTTTTTATAAACTGTTGTTACTCTTTAATTGACATTTATTTTTAAGCT
 TCCTAAGTTTGTGGTTCAAAAAAAAAAAAAAGGCGGCGTTACT

SEQ ID 17 = *Arabidopsis* obtusifoliol C14 α -
 demethylase(clone ID: ATA202967) amino acid translation

10 MELDSENKLLKTGLVIVATLVIKLIFFSFTSDSKKKRLPPTLKAWPPLVGSNIK
 FLKGPIIMLREEYPKLGSVFTVNLVHKKITFLIGPEVSAHFFKASESDLSSQEVY
 QFNVPTFGPGVVDVDYSVRQEQFRFFTEALRVNKLKGYVDMMVTEAEDYFSKWG
 ESGEVDIKVELERLIILTASRCLLGREVRDQLFDDVSALFHDLDNGMLPISVLF
 15 YLPIPAHRRDRAREKLSEIFAKIIGSRKRSKGTENDMLQCFIESKYKDGRQTTE
 SEVTGLLIAALFAGQHTSSITSTWTGAYLMRYKEYFSAALDEQKNLIAKHGDKID
 HDILSEMDVLYRCIKEALRLHPPLIMLRASHSDFSVTARDGKTYDIPKGHIVAT
 SPAFANRLPHIFKDPDTPERFSPGREEDKAAGAFSYIAFGGGRHGCLGEPFAY
 LQIKAIWSHLLRNFELELVSPFPEIDWNAMVVGKGNVMVRYKRRQLS*

20 SEQ ID 18 = *Arabidopsis* obtusifoliol C14 α -
 demethylase(clone ID: ATA403931) nucleotide sequence

TCGACCCCGCGTCCGCGGACGCGTGGGATCAGCTTCAAGCTTAAGAGAGC
 25 TTCGAAAGCGAAAGCGACGATTTCTTCTCCATCGTGAGAGCAAATCTCCAGAGCC
 GTTTTCTCTTCTTCTTCTTCTCCTCCTCGCGCCGTCTCTGAAACTCCATCATCGTAT
 CAATCAAATTGCTTCCCTCCTCAAATTGAAAAACAATGGAATTGGATTCCGAGAA
 CAAATTGTTGAAGACGGGTTTGGTTATAGTGGCGACACTTGTTATAGCCAAACTC
 ATCTTCTCTTTCTTCACTTCTGATTCTAAGAAGAAGCGTCTTCCCTCTACTCTTA
 30 AAGCTTGGCCTCCATTGGTTGGAAGTCTTATCAAATTCTTGAAAGGACCTATTAT
 TATGCTTAGAGAGGAATACCCTAAGCTTGGAAAGTGTGTTTACTGTTAATCTTGTT
 CACAAAAAGATTACTTTTCTTATTGGTCTGAAGTCTCTGCTCATTTTTTCAAAG
 CTTCTGAATCTGATCTTAGTCAGCAGGAAGTGTATCAGTTCAATGTCCCTACTTT
 TGGTCTCGGAGTTGTTTTCGATGTTGATTATTCTGTTCTGTCAGGAGCAGTTTCGG
 35 TTCTTCACTGAGGCACTTAGAGTTAACAAGTTGAAGGGTTATGTGGATATGATGG
 TTAAGCTGAGGCTGAGGATTACTTCTCTAAATGGGGAGAGAGTGGTGAAGTTGATAT
 TAAGGTTGAGCTAGAGAGGCTCATCATCTTGACTGCAAGTAGATGTTTACTGGGT
 CGAGAAGTTCGTGATCAGCTTTTGTATGATGTCTCTGCTTTGTTCCATGACCTTG
 ACAATGGAATGCTTCCCATCAGTGTTCTCTTCCCATATCTCCCAATTCCAGCTCA
 40 CCGCCGTGCTGACCGTGCCCGAGAAAAGCTTTCGGAGATTTTCGCAAAAATCATT
 GGGTCGAGAAAACGCTCTGGA AAAACAGAGAACGACATGCTGCAGTGTTCATCG
 AATCAAAGTACAAAGATGGTAGACAGACAACCGAATCTGAAGTCACTGGTTTGCT
 CATTGCTGCTCTGTtTGCAGGACAACACACGAGCTCTATCACTTCCACCTGGACC
 GGTGCTTATCTGATGCGATACAAAGAGTACTTCTCAGCTGCTCTTGATGAGCAGA
 45 AGAACCTGATTGCGAAACATGGAGACAAGATCGATCATGATATCTTATCCGAGAT
 GGATGTTCTCTACCGCTGCATTAAGGAAGCGTTGAGGCTTCACCCTCCACTCATC
 ATGTTAATGAGAGCCTCGCACAGTGATTTTACGCGTGACAGCTCGGGATGGAAAAA
 CTTACGATATCCCAAAGGGTCACATCGTTGCAACCTCCCCTGCATTTGCCAACCG
 CTTACCGCACATCTTCAAAGACCCCGACACCTACGACCCAGAAAGATTCTCCCT
 50 GGAAGAGAAGAGGACAAAGCCGCGAGGGGCATTCTCGTACATTGCATTCCGAGGGG

GAAGGCACGGGTGCCTTGGAGAGCCGTTTGCTTACCTGCAGATCAAAGCCATATG
GAGTCATTTGTTGAGGAACCTTCGAGCTTGAGCTAGTTTCACCGTTCCTGAGATT
GACTGGAACGCTATGGTGGTTGGAGTTAAAGGCAATGTGATGGTGCGTTACAAGA
GGCgCAGCTTTCTTAAAGACAAGTTTAAAGGTTATTGCAGCTTTGGATTTTCTC
5 TCTGGTTTCTGCTTTGCTTTTGTCCCTCTCTGGTTTTAGTTTTGTTGTTGAATAA
TTCTTCTGTTTTTATAAACTGTTGTTACTCTTTAATTGACATTTATTTTAAAGCT
TCCTAAGTTTGTGGTTCAAAAAAAAAAAAAAGGCGGCGTTACT

SEQ ID 19 = *Arabidopsis* obtusifoliol C14 α -
10 demethylase(clone ID: ATA403931) amino acid translation

MELDSENKLLKTGLVIVATLVIKLIFFSTSDSKKKRLPPTLKAWPPLVGSLIK
FLKGPIIMLREEYPKLGSVFTVNLVHKKITFLIGPEVSAHFFKASEDLSQQEVY
QFNVPFTFGPGVFDVDYSVRQEQFRFFTEALRVNKLKGYVDMMVTEAEDYFSKWG
15 ESGEVDIKVELERLIILTASRCLLGREVRDQLFDDVSALFHDLDNGMLPISVLF
YLPPIAHRRRDRAREKLSEIFAKIIGSRKRSKGTENDMLQCFIESKYKDGRTTE
SEVTGLLIAALFAGQHTSSITSTWTGAYLMRYKEYFSAALDEQKNLIAKHGDKID
HDILSEMDVLYRCIKEALRLHPPILMLMRASHSDFSVTARDGKTYDIPKGHIVAT
SPAFAANRLPHIFKDPDTPYDPERFSPGREEDKAAGAFSYIAFGGGRHGCLGEPFAY
20 LQIKAIWSHLLRNFELELVSPFPEIDWNAMVVGKGNVMVRYKRRQLS*

SEQ ID 20 = *Arabidopsis* sterol methyl transferase
I protein sequence (Accession NO: U71400) **See Appendix**
D

25 SEQ ID 21 = Tobacco sterol methyl transferase I
protein sequence (from Prof. Pierre Benveniste
Accession NO: U81312) **See Appendix E**

30 SEQ ID 22 = *Arabidopsis* sterol methyl transferase
II protein sequence (Accession NO: X89867) (from Prof.
Pierre Benveniste) **See Appendix F**

SEQ ID 23 = *Arabidopsis* sterol C5-desaturase
35 protein sequence (Accession NO: X90454) **See Appendix G**

SEQ ID 24 = Rubber truncated HMGR1m1 (S566 to A)
nucleotide sequence
ATGGCAGCGCCTCCCATGACGTGTGGGACCTCGAAGATACGGATCCCAACTACC
40 TCATCGATGAAGATCACCGTCTCGTTACTTGCCCTCCCGCTAATATATCTACTAA
GACTACCATTATTGCCGCACCTACCAAATTGCCTACCTCGGAACCCTTAATTGCA
CCCTTAGTCTCGGAGGAAGACGAAATGATCGTCAACTCCGTCGTGGATGGGAAGA
TACCTCCTATTCTCTGGAGTCGAAGCTCGGGGACTGCAAACGAGCGGCTGCGAT
TCGACGCGAGGCTTTGCAGAGGATGACAAGGAGGTGCTGGAAGGCTTGCCAGTA
45 GAAGGGTTCGATTACGAGTCGATTTTAGGACAATGCTGTGAAATGCCAGTGGGAT
ACGTGCAGATTCCGGTGGGGATTGCGGGGCCGTTGTTGCTGAACGGGCGGGAGTA
CTCTGTTCCAATGGCGACCACGGAGGGTTGTTTGGTGGCGAGCACTAATAGAGGG
TGTAAGGCGATTTACTTGTGAGGTGGGGCCACCAGCGTCTTGTTGAAGGATGGCA
TGACAAGAGCGCCTGTTGTAAGATTGCGGTCGGCGACTAGAGCCGCGGAGTTGAA
50 GTTCTTCTTGGAGGATCCTGACAATTTTGATACCTTGCCGTAAGTTTAAACAAG

098573 062004
T00290 2245860

TCCAGTAGATTTGCGAGGCTCCAAGGCATTAAATGCTCAATTGCTGGTAAGAATC
TTTATATAAGATTCAGCTGCAGCACTGGCGATGCAATGGGGATGAACATGGTTTC
TAAAGGGGTTCAAAACGTTCTTGAATTTCTTCAAAGTGATTTTTCTGATATGGAT
GTCATTGGAATCTCAGGAAATTTTTGTTTCGGATAAGAAGCCTGCTGCTGTAAATT
5 GGATTGAAGGACGTGGCAAATCAGTTGTTTGTGAGGCAATTATCAAGGAAGAGGT
GGTGAAGAAGGTGTTGAAAACCAATGTGGCCTCCCTAGTGGAGCTTAACATGCTC
AAGAATCTTGCTGGTTCTGCTGTTGCTGGTGCTTTGGGTGGATTTAATGCCCATG
CAGGCAACATCGTATCTGCAATCTTTATTGCCACTGGCCAGGATCCAGCACAGAA
TGTTGAGAGTTCTCATTGCATTACCATGATGGAAGCTGTCAATGATGGAAGGAT
10 CTCCATATCTCTGTGACCATGCCCTCCATTGAGGTGGGTACAGTCGGAGGTGGAA
CTCAACTTGCATCTCAGTCTGCTTGTCTCAATTTGCTTGGGGTGAAGGGTGCAAA
CAAAGAGTCGCCAGGATCAAACCTCAAGGCTCCTTGCTGCCATCGTAGCTGGTTCA
GTTTTGGCTGGTGAGCTCTCCTTGATGTCTGCCATTGCAGCTGGGCAGCTTGTC
AGAGTCACATGAAGTACAACAGAGCCAGCAAAGATATGTCTAAAGCTGCATCTTA
15 G

SEQ ID 25 = Rubber truncated HMGR1m1 (S566 to A)
amino acid translation

MARASHDVWDLEDTPNYLIDEDHRLVTCPPANISTKTTIIAAPTCLPTSEPLIA
20 PLVSEEDEMIVNSVVDGKIPSYSLKLGDKRAAAIRREALQRMTRRSLEGLPV
EGFDYESILGQCCMPVGYVQIPVGIAGPLLLNGREYSVPMATTEGCLVASTNRG
CKAIYLSGGATSULLKDGMTAPVVRFASATRAAELKFFLEDPDNFDTLAVVFNK
SSRFARLQGIKCSIAGKNLYIRFSCSTGDAMGMNMVSKGVQNVLEFLQSDFSMD
VIGISGNFCSDDKPAAVNWIEGRGKSVVCEAIKKEEVVKKVLKTNVASLVELNML
25 KNLAGSAVAGALGGFNAHAGNIVSAIFIATGQDPAQNVESHCITMMEAVNDGKD
LHISVTMPSIEVGTVGGGTQLASQSACLNLGKVGANKESPGSNSRLLAAIVAGS
VLAGELSLMSAIAAGQLVKSHMKYNRASKDMSKAAS

SEQ ID. 26 = Rubber truncated HMGR1m2 (S567 to A)
nucleotide sequence

ATGGCACGCGCTCCCATGACGTGTGGGACCTCGAAGATACGGATCCCAACTACC
TCATCGATGAAGATCACCGTCTCGTTACTTGCCCTCCCGCTAATATATCTACTAA
35 GACTACCATTATTGCCGCACCTACCAAATTGCCTACCTCGGAACCTTAATTGCA
CCCTTAGTCTCGGAGGAAGACGAAATGATCGTCAACTCCGTCGTGGATGGGAAGA
TACCCTCCTATTCTCTGGAGTCGAAGCTCGGGGACTGCAAACGAGCGGCTGCGAT
TCGACGCGAGGCTTTGCAGAGGATGACAAGGAGGTGCTGGAAGGCTTGCCAGTA
GAAGGGTTCGATTACGAGTCGATTTTAGGACAATGCTGTGAAATGCCAGTGGGAT
40 ACGTGCAGATTCCGGTGGGGATTGCGGGGCCGTTGTTGCTGAACGGGCGGGAGTA
CTCTGTTCCAATGGCGACCACGGAGGGTGTGTTGGTGGCGAGCACTAATAGAGGG
TGTAAGGCGATTACTTGTGAGGTGGGGCCACCAGCGTCTTGTTGAAGGATGGCA
TGACAAGAGCGCCTGTTGTAAGATTGCGGTGCGGACTAGAGCCGCGGAGTTGAA
GTTCTTCTTGAGAGATCCTGACAATTTTGATACCTTGCCGCTAGTTTTTAACAAG
45 TCCAGTAGATTTGCGAGGCTCCAAGGCATTAAATGCTCAATTGCTGGTAAGAATC
TTTATATAAGATTCAGCTGCAGCACTGGCGATGCAATGGGGATGAACATGGTTTC
TAAAGGGGTTCAAAACGTTCTTGAATTTCTTCAAAGTGATTTTTCTGATATGGAT
GTCATTGGAATCTCAGGAAATTTTTGTTTCGGATAAGAAGCCTGCTGCTGTAAATT
GGATTGAAGGACGTGGCAAATCAGTTGTTTGTGAGGCAATTATCAAGGAAGAGGT
50 GGTGAAGAAGGTGTTGAAAACCAATGTGGCCTCCCTAGTGGAGCTTAACATGCTC

09865723.062004

AAGAATCTTGCTGGTTCTGCTGTTGCTGGTGCTTTGGGTGGATTTAATGCCCATG
 CAGGCAACATCGTATCTGCAATCTTTATTGCCACTGGCCAGGATCCAGCACAGAA
 TGTGAGAGTTCTCATTGCATTACCATGATGGAAGCTGTCAATGATGGAAAGGAT
 CTCCATATCTCTGTGACCATGCCCTCCATTGAGGTGGGTACAGTCGGAGGTGGAA
 5 CTCAACTTGCATCTCAGTCTGCTTGTCTCAATTTGCTTGGGGTGAAGGTGCAAA
 CAAAGAGTCGCCAGGATCAAACAGGCTCCTTGCTGCCATCGTAGCTGGTTCA
 GTTTTGGCTGGTGAGCTCTCCTTGATGTCTGCCATTGCAGCTGGGCAGCTTGTC
 AGAGTCACATGAAGTACAACAGATCCGCCAAAGATATGTCTAAAGCTGCATCTTA
 G

10

SEQ ID 27 = Rubber truncated HMGR1m2 (S567 to A)
 amino acid translation
 MARASHDVWDLEDTPNYLIDEDHRLVTCPPANISTKTTIIAAPTCLPTSEPLIA
 15 PLVSEEDEMIVNSVVDGKIPSYSLKLGDKRAAAIRREALQRMTRRSLEGLPV
 EGFDESILGQCCMPVGYVQIPVGIAGPLLLNGREYSVPMATTEGCLVASTNRG
 CKAIYLSGGATSVLLKDGMTAPVVRFASATRAAELKFFLEDPDNFDTLAVVFNK
 SSRFARLQGIKCSIAGKNLYIRFSCSTGDAMGMNMVSKGVQNVLEFLQSDFSMD
 VIGISGNFCSDKKPAAVNWIEGRGKSVVCEAI IKEEVKKVLKTNVASLVELNML
 20 KNLAGSAVAGALGGFNAHAGNIVSAIFIATGQDPAQNVESHCHITMMEAVNDGKD
 LHSVTMPSEIEVGTVGGGTQLASQSACLNLLGVKGANKESPGSNSRLLAAIVAGS
 VLAGELSLMSAIAAGQLVKSHMKYNRSKDMSKAAS

Example 6. *Arabidopsis* obtusifoliol C14 α -
 25 demethylase constructs

The *Arabidopsis* obtusifoliol C14 α -demethylase gene
 was amplified from two separate *Arabidopsis* mRNA
 samples (SIN 2 and Keto-10) through use of primers
 BXK33 and BXK34, as described below.

30

BXK33 (SEQ ID 28): 5'-**GAGATCT**GAACCCTAACGAGAG-3'

BXK34 (SEQ ID 29): 5'-**GGAGCTCT**TAAAGAAAAGGGACGACGC-3'

35 The primer BXK33 has a Bgl II cleavage site shown
 in bold. The primer BXK34 has a Sac I cleavage site
 shown in bold. The actual size of the structural gene
 is 1.445 Kb.

The *Arabidopsis* mRNA was amplified using a Perkin
 40 Elmer GeneAmp RT-PCR kit. The reverse transcription
 reaction used 25 mM MgCl₂ (4 μ l; 5 mM final), 10X PCR

T00290" E25860

buffer (2 μ l), di DEPC water (1 μ l), 2 μ l each of 1 mM solution of each of four dNTPs (dGTP, dATP, dUTP, dCTP), RNase inhibitor (1 μ l of 10 units per μ l stock), MMLV reverse transcriptase (1 μ l of a 2.5 U/ μ l stock),
 5 Oligo d(T)16 Primer (1 μ l of a 2.5 μ M stock), and 2 μ l of an *Arabidopsis* polyA RNA sample. The reaction mix was incubated at room temperature (about 20°C) for 10 minutes, then in a PCR machine for one cycle (15 min. at 42°C, 5 min. at 99°C and 5 min. at 4°C).

10 Separate primer-mediated amplification reactions were carried out using Taq DNA polymerase and Vent DNA polymerase to obtain *Arabidopsis obtusifolius* Cl4 α -demethylase cDNA from the amplified mRNA sample.

Taq PCR Reaction	Vent PCR Reaction
4 μ l 25 mM MgCl ₂	4 μ l 25 mM MgCl ₂
8 μ l 10X PCR buffer II	8 μ l 10X Vent PCR buffer
65.5 μ l di DEPC water	65.5 μ l di DEPC water
0.5 μ l AmpliTaq polymerase	0.5 μ l Vent polymerase

15

After 1 minute and 35 seconds at 95°C, 1 μ l each of 15 μ M stocks of the upstream and downstream primers (BXK33 and BXK34) were added to the PCR reaction (100 μ l total PCR reaction volume) and the PCR reaction
 20 solutions were subjected to 35 cycles (95°C for 15 seconds, then 60°C for 30 seconds). The amplified PCR reaction was then maintained at 72°C for 7 minutes and then stored at 4°C. An amplification positive control reaction was carried out under the same conditions with
 25 DM151 and DM152 primers.

DM151 (SEQ ID 30): 5'-GTCTCTGAATCAGAAATCCTTCTATC-3'

100290" E2258860

DM152 (SEQ ID 31): 5'-CATGTCAAATTTCACTGCTTCATCC-3'

Electrophoresis of the nucleic acid solutions after PCR amplification displayed an amplification product corresponding approximately to the size of the desired 1.445 Kb structural gene. The fragment was cloned into an M13 vector. A representative sequencing reaction consisted of: 10 µl of plasmid DNA (200-500 ng), 2 µl of M13 Forward or Reverse primer (15 picomoles) and 8 µl of Big Dye Terminator Reaction Mix (PE Applied Biosci.). The clone copy of ATA101105 was called CPR17398. The sequence of the selected clone (*Arabidopsis obtusifoliol* C14α-demethylase) is identified as SEQ ID NO:9.

The predicted polypeptide sequence for the cloned *Arabidopsis obtusifoliol* C14α-demethylase sequence was subjected to a BLAST search in the public database and found to align with the sorghum obtusifoliol 14-alpha demethylase polypeptide (ATA101105/U74319/g1658192; and g1216657/U74319) exhibiting 75-78% sequence identity and 87-90% sequence homology. The cloned nucleic acid encoding *Arabidopsis obtusifoliol* C14α-demethylase (SEQ ID No:9) is missing the 5' end.

The 5' terminal portion of the structural gene was obtained by the RACE (Rapid Amplification of cDNA Ends) PCR using primers BXK39 and BXK40 per manufacturer's instructions (Clontech).

BXK39 (SEQ ID 32): 5'-GAGATCTCCACAGATTTAAAGAACCCTAACG-3'

BXK40 (SEQ ID 33): 5'-GGAGCTCGGTTTTTAAGAAAAGGGACGACGC-3'

The cloned nucleic acid encoding full length *Arabidopsis obtusifoliol* C14α-demethylase is identified

as SEQ ID No:8. The amplified *Arabidopsis obtusifoliol* C14 α -demethylase structural gene is useful for making constructs that express *Arabidopsis obtusifoliol* C14 α -demethylase in transgenic plants.

5 In light of the detailed description of the invention and the examples presented above, it can be appreciated that the several aspects of the invention are achieved.

It is to be understood that the present invention
10 has been described in detail by way of illustration and example in order to acquaint others skilled in the art with the invention, its principles, and its practical application. Particular formulations and processes of the present invention are not limited to the
15 descriptions of the specific embodiments presented, but rather the descriptions and examples should be viewed in terms of the claims that follow and their equivalents. While some of the examples and descriptions above include some conclusions about the
20 way the invention may function, the inventor does not intend to be bound by those conclusions and functions, but puts them forth only as possible explanations.

It is to be further understood that the specific
embodiments of the present invention as set forth are
25 not intended as being exhaustive or limiting of the invention, and that many alternatives, modifications, and variations will be apparent to those of ordinary skill in the art in light of the foregoing examples and detailed description. Accordingly, this invention is
30 intended to embrace all such alternatives, modifications, and variations that fall within the spirit and scope of the following claims.

09885729.062091
"E245860"